

Reimbursement Policy

Fecal Calprotectin Testing in Adults

[POLICY DESCRIPTION](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) |

I. Policy Description

Calprotectin is a small calcium-binding protein found in high concentration in the cytosol of neutrophils (Fagerhol et al., 1980) and to a lesser extent monocytes and macrophages (Hsu et al., 2009). Active intestinal inflammation and disturbance of the mucosa results in entrance of neutrophils (containing calprotectin) into the lumen and subsequent excretion in feces. Detection of fecal calprotectin is used to distinguish inflammatory bowel disease (IBD) from irritable bowel syndrome (IBS) and other causes of abdominal discomfort, bloating, and diarrhea (Walsham & Sherwood, 2016).

II. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the "Applicable State and Federal Regulations" section of this policy document.

This policy is specific to individuals 18 years of age or older. Criteria below do not apply to individuals less than 18 years of age.

- 1) Fecal calprotectin testing for the differential diagnosis between non-inflammatory gastrointestinal disease (e.g., IBS) and inflammatory gastrointestinal disease (e.g., IBD) **MEETS COVERAGE CRITERIA.**
- 2) Fecal calprotectin testing either to assess for response to therapy or for relapse or to monitor gastrointestinal conditions such as inflammatory bowel disease (IBD) **MEETS COVERAGE CRITERIA.**

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.

- 3) Fecal calprotectin testing for all other situations not discussed above **DOES NOT MEET COVERAGE CRITERIA.**

III. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this Policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National

Reimbursement Policy

Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <https://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

In March 2006, the PhiCal™ (Genova Diagnostics) quantitative ELISA test for measuring concentrations of fecal calprotectin in fecal stool was cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) processes. This test is indicated to aid in the diagnosis of inflammatory bowel disease (IBD) and to differentiate IBD from irritable bowel syndrome (IBS); it is intended to be used in conjunction with other diagnostic testing and clinical considerations (FDA, 2006). On December 26, 2018, a successor device called “LIAISON Calprotectin, LIAISON Calprotectin Control Set, LIAISON Calprotectin Calibration Verifiers, LIAISON Q.S.E.T. Buffer, LIAISON Q.S.E.T. Device” was approved. The new description is as follows: “The DiaSorin LIAISON® Calprotectin assay is an in vitro diagnostic chemiluminescent immunoassay (CLIA) intended for the quantitative measurement, in human stool, of fecal calprotectin, a neutrophilic protein that is a marker of mucosal inflammation. The LIAISON® Calprotectin assay can be used as an aid in the diagnosis of inflammatory bowel diseases (IBD), specifically Crohn’s disease and ulcerative colitis, and as an aid in differentiation of IBD from irritable bowel syndrome (IBS). Test results are to be used in conjunction with information obtained from the patients’ clinical evaluation and other diagnostic procedures. The test has to be performed on the LIAISON® XL Analyzer” (FDA, 2018a).

In January 2014, CalPrest® (Eurospital SpA, Trieste, Italy) was cleared for marketing by the FDA through the 510(k) processes. According to the FDA summary, CalPrest® “is identical” to the PhiCal™ test “in that they are manufactured by Eurospital S.p.A. Trieste, Italy. The only differences are the name of the test on the labels, the number of calibrators in the kit and the dynamic range of the assay.” CalPrest®NG (Eurospital SpA) was cleared for marketing in November 2016 (FDA, 2016).

On October 16, 2018, the FDA approved the QUANTA Flash Calprotectin and Fecal Extraction Device. The device’s intended use is as follows: “QUANTA Flash Calprotectin is a chemiluminescent immunoassay for the quantitative determination of fecal calprotectin in extracted human stool samples. Elevated levels of fecal calprotectin, in conjunction with clinical findings and other laboratory tests, can aid in the diagnosis of inflammatory bowel disease (IBD) (ulcerative colitis and Crohn’s disease), and in the differentiation of IBD from irritable bowel syndrome (IBS).” This device has a predicate device, which was approved in 2017 (FDA, 2018a).

On December 26, 2018, the FDA approved the LIAISON Calprotectin Assay. The device’s intended use is as follows: “The DiaSorin LIAISON® Calprotectin assay is an in vitro diagnostic

Reimbursement Policy

chemiluminescent immunoassay (CLIA) intended for the quantitative measurement, in human stool, of fecal calprotectin, a neutrophilic protein that is a marker of mucosal inflammation. The LIAISON® Calprotectin assay can be used as an aid in the diagnosis of inflammatory bowel diseases (IBD), specifically Crohn’s disease and ulcerative colitis, and as an aid in differentiation of IBD from irritable bowel syndrome (IBS). Test results are to be used in conjunction with information obtained from the patients’ clinical evaluation and other diagnostic procedures” (FDA, 2018b).

On September 24, 2019, BÜHLMANN Laboratories AG received FDA approval for the Buhlmann FCAL Turbo and CALEX Cap fecal calprotectin extraction device. This device is to be used in conjunction with the automated calprotectin test, BÜHLMANN fCAL® turbo. The BÜHLMANN fCAL® turbo is an in vitro diagnostic assay which quantitatively measures fecal calprotectin (FDA, 2019).

Rapid fecal calprotectin tests, such as CalproSmart™, are available internationally for use as point-of-care testing, but these have not been approved for use in the U.S. by the FDA.

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

IV. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
83993	Calprotectin, fecal

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

V. Evidence-based Scientific References

- Abej, E., El-Matary, W., Singh, H., & Bernstein, C. N. (2016). The Utility of Fecal Calprotectin in the Real-World Clinical Care of Patients with Inflammatory Bowel Disease. *Can J Gastroenterol Hepatol*, 2016, 2483261. <https://doi.org/10.1155/2016/2483261>
- Almario, C. V., Ballal, M. L., Chey, W. D., Nordstrom, C., Khanna, D., & Spiegel, B. M. R. (2018). Burden of Gastrointestinal Symptoms in the United States: Results of a Nationally Representative Survey of Over 71,000 Americans. *Am J Gastroenterol*, 113(11), 1701-1710. <https://doi.org/10.1038/s41395-018-0256-8>
- Boirivant, M., & Cossu, A. (2012). Inflammatory bowel disease. *Oral Dis*, 18(1), 1-15. <https://doi.org/10.1111/j.1601-0825.2011.01811.x>

Reimbursement Policy

- Burri, E., & Beglinger, C. (2014). The use of fecal calprotectin as a biomarker in gastrointestinal disease. *Expert Rev Gastroenterol Hepatol*, 8(2), 197-210. <https://doi.org/10.1586/17474124.2014.869476>
- Campbell, J. P., Zierold, C., Rode, A. M., Blocki, F. A., & Vaughn, B. P. (2021). Clinical Performance of a Novel LIAISON Fecal Calprotectin Assay for Differentiation of Inflammatory Bowel Disease From Irritable Bowel Syndrome. *J Clin Gastroenterol*, 55(3), 239-243. <https://doi.org/10.1097/mcg.0000000000001359>
- Colombel, J. F., Shin, A., & Gibson, P. R. (2019). AGA Clinical Practice Update on Functional Gastrointestinal Symptoms in Patients With Inflammatory Bowel Disease: Expert Review. *Clin Gastroenterol Hepatol*, 17(3), 380-390.e381. <https://doi.org/10.1016/j.cgh.2018.08.001>
- Fagerhol, M. K., Dale, I., & Andersson, T. (1980). A radioimmunoassay for a granulocyte protein as a marker in studies on the turnover of such cells. *Bull Eur Physiopathol Respir*, 16 Suppl, 273-282.
- FDA. (2006). 510(k) SUBSTANTIAL EQUIVALENCE DETERMINATION https://www.accessdata.fda.gov/cdrh_docs/reviews/K050007.pdf
- FDA. (2016). 510(k) https://www.accessdata.fda.gov/cdrh_docs/pdf16/K160447.pdf
- FDA. (2018a). 510(k) https://www.accessdata.fda.gov/cdrh_docs/pdf18/K182698.pdf
- FDA. (2018b). LIAISON Calprotectin. https://www.accessdata.fda.gov/cdrh_docs/pdf18/K182698.pdf
- FDA. (2019). *Buhlmann FCAL Turbo And CALEX Cap.* <https://www.accessdata.fda.gov/scripts/cdrh/devicesatfda/index.cfm?db=pmn&id=K191718>
- Gibson, P. (2022). *Irritable bowel syndrome in patients with inflammatory bowel disease - UpToDate.* <https://www.uptodate.com/contents/irritable-bowel-syndrome-in-patients-with-inflammatory-bowel-disease>
- Gomollón, F., Dignass, A., Annesse, V., Tilg, H., Van Assche, G., Lindsay, J. O., Peyrin-Biroulet, L., Cullen, G. J., Daperno, M., Kucharzik, T., Rieder, F., Almer, S., Armuzzi, A., Harbord, M., Langhorst, J., Sans, M., Chowers, Y., Fiorino, G., Juillerat, P., . . . on behalf of, E. (2016). 3rd European Evidence-based Consensus on the Diagnosis and Management of Crohn's Disease 2016: Part 1: Diagnosis and Medical Management. *Journal of Crohn's and Colitis*, 11(1), 3-25. <https://doi.org/10.1093/ecco-jcc/jjw168>
- Halpin, S. J., & Ford, A. C. (2012). Prevalence of symptoms meeting criteria for irritable bowel syndrome in inflammatory bowel disease: systematic review and meta-analysis. *Am J Gastroenterol*, 107(10), 1474-1482. <https://doi.org/10.1038/ajg.2012.260>
- Higuchi, L. M., & Bousvaros, A. (2024). Clinical presentation and diagnosis of inflammatory bowel disease in children - UpToDate. In M. Heyman (Ed.), *UpToDate.* <https://www.uptodate.com/contents/clinical-presentation-and-diagnosis-of-inflammatory-bowel-disease-in-children>
- Hsu, K., Champai boon, C., Guenther, B. D., Sorenson, B. S., Khammanivong, A., Ross, K. F., Geczy, C. L., & Herzberg, M. C. (2009). ANTI-INFECTIVE PROTECTIVE PROPERTIES OF S100 CALGRANULINS. *Antiinflamm Antiallergy Agents Med Chem*, 8(4), 290-305. <https://doi.org/10.2174/187152309789838975>

Reimbursement Policy

- Johnson, L. M., Spannagl, M., Wojtalewicz, N., & Durner, J. (2022). Comparison of fecal calprotectin and pancreatic elastase assays based on proficiency testing results. *Clin Biochem*, *107*, 19-23. <https://doi.org/10.1016/j.clinbiochem.2022.05.002>
- Khaki-Khatibi, F., Qujeq, D., Kashifard, M., Moein, S., Maniati, M., & Vaghari-Tabari, M. (2020). Calprotectin in inflammatory bowel disease. *Clin Chim Acta*, *510*, 556-565. <https://doi.org/10.1016/j.cca.2020.08.025>
- Lacy, B. E., Pimentel, M., Brenner, D. M., Chey, W. D., Keefer, L. A., Long, M. D., & Moshiree, B. (2021). ACG Clinical Guideline: Management of Irritable Bowel Syndrome. *Am J Gastroenterol*, *116*(1), 17-44. <https://doi.org/10.14309/ajg.0000000000001036>
- Lichtenstein, G. R., Loftus, E. V., Isaacs, K. L., Regueiro, M. D., Gerson, L. B., & Sands, B. E. (2018). ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol*, *113*(4), 481-517. <https://doi.org/10.1038/ajg.2018.27>
- Maaser, C., Sturm, A., Vavricka, S. R., Kucharzik, T., Fiorino, G., Annese, V., Calabrese, E., Baumgart, D. C., Bettenworth, D., Borralho Nunes, P., Burisch, J., Castiglione, F., Eliakim, R., Ellul, P., González-Lama, Y., Gordon, H., Halligan, S., Katsanos, K., Kopylov, U., . . . Stoker, J. (2019). ECCO-ESGAR Guideline for Diagnostic Assessment in IBD Part 1: Initial diagnosis, monitoring of known IBD, detection of complications. *J Crohns Colitis*, *13*(2), 144-164. <https://doi.org/10.1093/ecco-jcc/jjy113>
- Magro, F., Gionchetti, P., Eliakim, R., Ardizzone, S., Armuzzi, A., Barreiro-de Acosta, M., Burisch, J., Gecse, K. B., Hart, A. L., Hindryckx, P., Langner, C., Limdi, J. K., Pellino, G., Zagórowicz, E., Raine, T., Harbord, M., Rieder, F., for the European, C. s., & Colitis, O. (2017). Third European Evidence-based Consensus on Diagnosis and Management of Ulcerative Colitis. Part 1: Definitions, Diagnosis, Extra-intestinal Manifestations, Pregnancy, Cancer Surveillance, Surgery, and Ileo-anal Pouch Disorders. *Journal of Crohn's and Colitis*, *11*(6), 649-670. <https://doi.org/10.1093/ecco-jcc/jjx008>
- Mao, R., Xiao, Y. L., Gao, X., Chen, B. L., He, Y., Yang, L., Hu, P. J., & Chen, M. H. (2012). Fecal calprotectin in predicting relapse of inflammatory bowel diseases: a meta-analysis of prospective studies. *Inflamm Bowel Dis*, *18*(10), 1894-1899. <https://doi.org/10.1002/ibd.22861>
- Molander, P., af Bjorkesten, C. G., Mustonen, H., Haapamaki, J., Vauhkonen, M., Kolho, K. L., Farkkila, M., & Sipponen, T. (2012). Fecal calprotectin concentration predicts outcome in inflammatory bowel disease after induction therapy with TNFalpha blocking agents. *Inflamm Bowel Dis*, *18*(11), 2011-2017. <https://doi.org/10.1002/ibd.22863>
- Molander, P., Farkkila, M., Ristimaki, A., Salminen, K., Kempainen, H., Blomster, T., Koskela, R., Jussila, A., Rautiainen, H., Nissinen, M., Haapamaki, J., Arkkila, P., Nieminen, U., Kuisma, J., Punkkinen, J., Kolho, K. L., Mustonen, H., & Sipponen, T. (2015). Does fecal calprotectin predict short-term relapse after stopping TNFalpha-blocking agents in inflammatory bowel disease patients in deep remission? *J Crohns Colitis*, *9*(1), 33-40. <https://doi.org/10.1016/j.crohns.2014.06.012>
- Mumolo, M. G., Bertani, L., Ceccarelli, L., Laino, G., Di Fluri, G., Albano, E., Tapete, G., & Costa, F. (2018). From bench to bedside: Fecal calprotectin in inflammatory bowel diseases

Reimbursement Policy

- clinical setting. *World J Gastroenterol*, 24(33), 3681-3694. <https://doi.org/10.3748/wjg.v24.i33.3681>
- NICE. (2017). Faecal calprotectin diagnostic tests for inflammatory diseases of the bowel DG11. *NICE Diagnostics guidance*. <https://www.nice.org.uk/guidance/DG11>
- Rosenfeld, G., Greenup, A. J., Round, A., Takach, O., Halparin, L., Saadeddin, A., Ho, J. K., Lee, T., Enns, R., & Bressler, B. (2016). FOCUS: Future of fecal calprotectin utility study in inflammatory bowel disease. *World J Gastroenterol*, 22(36), 8211-8218. <https://doi.org/10.3748/wjg.v22.i36.8211>
- Rubin, D. T., Ananthakrishnan, A. N., Siegel, C. A., Sauer, B. G., & Long, M. D. (2019). ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol*, 114(3), 384-413. <https://doi.org/10.14309/ajg.0000000000000152>
- Singh, S., Ananthakrishnan, A. N., Nguyen, N. H., Cohen, B. L., Velayos, F. S., Weiss, J. M., Sultan, S., Siddique, S. M., Adler, J., & Chachu, K. A. (2023). AGA Clinical Practice Guideline on the Role of Biomarkers for the Management of Ulcerative Colitis. *Gastroenterology*, 164(3), 344-372. <https://doi.org/10.1053/j.gastro.2022.12.007>
- Tham, Y. S., Yung, D. E., Fay, S., Yamamoto, T., Ben-Horin, S., Eliakim, R., Koulaouzidis, A., & Kopylov, U. (2018). Fecal calprotectin for detection of postoperative endoscopic recurrence in Crohn's disease: systematic review and meta-analysis. *Therap Adv Gastroenterol*, 11, 1756284818785571. <https://doi.org/10.1177/1756284818785571>
- Tibble, J. A., Sigthorsson, G., Foster, R., Forgacs, I., & Bjarnason, I. (2002). Use of surrogate markers of inflammation and Rome criteria to distinguish organic from nonorganic intestinal disease. *Gastroenterology*, 123(2), 450-460. <https://doi.org/10.1053/gast.2002.34755>
- van Rheenen, P. F., Van de Vijver, E., & Fidler, V. (2010). Faecal calprotectin for screening of patients with suspected inflammatory bowel disease: diagnostic meta-analysis. *Bmj*, 341, c3369. <https://doi.org/10.1136/bmj.c3369>
- von Roon, A. C., Karamountzos, L., Purkayastha, S., Reese, G. E., Darzi, A. W., Teare, J. P., Paraskeva, P., & Tekkis, P. P. (2007). Diagnostic precision of fecal calprotectin for inflammatory bowel disease and colorectal malignancy. *Am J Gastroenterol*, 102(4), 803-813. <https://doi.org/10.1111/j.1572-0241.2007.01126.x>
- Walsham, N. E., & Sherwood, R. A. (2016). Fecal calprotectin in inflammatory bowel disease. *Clin Exp Gastroenterol*, 9, 21-29. <https://doi.org/10.2147/ceg.s51902>
- Wagh, N., Cummins, E., Royle, P., Kandala, N. B., Shyangdan, D., Arasaradnam, R., Clar, C., & Johnston, R. (2013). Faecal calprotectin testing for differentiating amongst inflammatory and non-inflammatory bowel diseases: systematic review and economic evaluation. *Health Technol Assess*, 17(55), xv-xix, 1-211. <https://doi.org/10.3310/hta17550>
- Yang, Z., Clark, N., & Park, K. T. (2014). Effectiveness and cost-effectiveness of measuring fecal calprotectin in diagnosis of inflammatory bowel disease in adults and children. *Clin Gastroenterol Hepatol*, 12(2), 253-262.e252. <https://doi.org/10.1016/j.cgh.2013.06.028>