

Reimbursement Policy

Prescription Medication and Illicit Drug Testing in the Outpatient Setting

[POLICY DESCRIPTION](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [REIMBURSEMENT](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) | [REVISION HISTORY](#)

I. Policy Description

Abuse of both prescription and illicit drugs is extremely common. Drugs of abuse (DOA) may be defined as “a drug, chemical, or plant product that is known to be misused for recreational purposes,” which can include drugs such as pain relievers that have legitimate prescriptions. Drug tests may be performed for a variety of reasons, such as compliance with treatment program or medical regimen. Numerous biological substances, such as blood, hair, or saliva may be tested, but urine is the most commonly tested biological substance in drug tests (Hoffman, 2023).

This policy addresses clinical toxicology in the outpatient setting and does not address forensic testing or therapeutic drug monitoring (TDM). Forensic drug testing is used for legal proceedings and requires secondary confirmatory testing (Jones, 2016). TDM “involves sampling of plasma or serum drug levels to determine optimal drug dosing” (Eaton & Lyman, 2022).

II. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual’s benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the “Applicable State and Federal Regulations” section of this policy document.

This policy concerns only coverage criteria and does not describe or define the legal responsibility of providers. Providers should refer to state and federal laws for such guidance.

This policy does not address the use of drug testing in the following circumstances:

- A. State, federally regulated, and legally mandated drug testing (i.e., court-ordered drug screening, forensic examinations).*
- B. Non-forensic testing for commercial driver’s licensing or any other job-related testing (i.e., as a prerequisite for employment or as a means for continuation of employment).*
- C. As a component of routine physical/medical examination.*
- D. As a component of care rendered in an urgent/emergency situation.*
- E. As a routine component of a behavioral health assessment.*

PRESUMPTIVE DRUG SCREENING USING URINE SAMPLES

Reimbursement Policy

- 1) Presumptive drug screening using urine samples (qualitative, semi-quantitative or quantitative) **MEETS COVERAGE CRITERIA** in **any** of the following situations:
 - a) To assess an individual being treated for chronic, non-cancer pain when clinical evaluation of the individual (history/signs/symptoms) suggests the use of non-prescribed medications or illegal substances:
 - i) Prior to initiating chronic opioid pain therapy in chronic non-cancer pain to determine if the individual has been exposed to controlled substances or potentially confounding illicit drugs.
 - ii) To verify an individual's compliance with treatment or identify undisclosed drug abuse as part of routine monitoring for individuals who are receiving treatment for non-cancer chronic pain with prescription opioid pain medication. The random testing interval and drugs selected for testing should be based on the individual's history, condition, and treatment, as documented in the medical record.
 - (a) Monitoring of low risk (as defined by a risk assessment tool) individuals on chronic opioid therapy, up to one (2) time per year after initiation of therapy.
 - (b) Monitoring of moderate risk (as defined by a risk assessment tool) individuals on chronic opioid therapy, up to two (4) times per year after initiation of therapy.
 - (c) Monitoring of high risk (as defined by a risk assessment tool) individuals on chronic opioid therapy, up to four (5) times per year after initiation of therapy.
 - (d) For individuals with aberrant behavior (lost prescriptions, multiple requests for early refills, and opioids from multiple providers, unauthorized dose escalation, apparent intoxication, etc.), testing at the time of visit meets coverage criteria.
 - b) In pregnant individuals at high-risk for substance abuse in whom the suspicion of drug use exists based on the answers to substance abuse screening questions or as indicated by information from the prescription drug monitoring program (PDMP), as documented in the medical record.
 - c) In newborns when there is a history of maternal substance abuse or agitated/altered mental status in the birthing parent.
 - d) In candidates for organ transplant who have a history of substance abuse (to demonstrate abstinence prior to transplant).

Reimbursement Policy

- e) In individuals with a suspicion of or a diagnosis of mental illness (e.g., anxiety disorders, schizophrenia, major depressive disorder, mood disorders, suicidal ideations, substance abuse disorder).
- f) In individuals with attention-deficit hyperactivity and disruptive behavior disorders.
- g) In cancer patients on opioid pain medication.
- h) In individuals with epilepsy.
- i) For the management and compliance monitoring of an individual under treatment for substance abuse or dependence at the following frequency (after baseline at initial evaluation) and must be documented in the patient's medical record:
 - i) For patients with 0 to 90 consecutive days of abstinence, random qualitative drug testing at a frequency of 1 to 2 per week.
 - ii) For patients with > 90 consecutive days of abstinence, random qualitative drug testing at a frequency of 1 to 3 per month.
- j) In individuals where substance abuse is in the differential diagnosis of the presenting conditions.

DEFINITIVE DRUG TESTING

- 2) Confirmatory/definitive qualitative or quantitative drug testing (up to seven drug classes) **MEETS COVERAGE CRITERIA** when laboratory-based definitive drug testing is specifically requested, the rationale is documented by the patient's treating physician, **and any** of the following conditions are met:
 - a) The result of the presumptive drug screen is different than that suggested by the patient's medical history, their clinical presentation, or patient's own statement (e.g., test was negative for prescribed medications, test was positive for prescription drug with abuse potential, which was not prescribed, test was positive for an illegal drug).
 - b) For diagnosing and monitoring individuals with substance use disorder or dependence, when accurate and reliable results are necessary for treatment decisions:
 - i) Individuals with 0 to 30 consecutive days of abstinence, random definitive drug testing at a frequency not to exceed 1 per week.
 - ii) Individuals with 31 to 90 consecutive days of abstinence, random definitive drug testing at a frequency of 1 to 3 per month. No more than 3 definitive drug tests in one month will be allowed.

Reimbursement Policy

- iii) Individuals with greater than 90 consecutive days of abstinence, definitive drug testing at a frequency of 1 to 3 every three months. No more than 3 definitive drug tests in a 3-month period will be allowed.
 - c) For monitoring of individuals on opioid therapy (to ensure adherence to the therapeutic plan, for treatment planning, and for detection of other, non-prescribed opioids).
 - d) A presumptive test does not exist or does not adequately detect the specific drug or metabolite to be tested (e.g., specific drugs within the amphetamine, barbiturate, benzodiazepine, tricyclic antidepressants, and opiate/opioid drug classes, as well as synthetic/analog or “designer” drugs).
 - e) To definitively identify specific drugs in a large family of drugs.
 - f) To identify drugs when a definitive concentration of a drug is needed to guide management.
- 3) When laboratory-based definitive drug testing is requested for larger than seven drug classes panels, confirmatory/definitive qualitative or quantitative drug testing **DOES NOT MEET COVERAGE CRITERIA**.
 - 4) Confirmatory/definitive qualitative or quantitative **or** presumptive (qualitative, semi-quantitative or quantitative) drug testing using proprietary tests (e.g., CareView360) **DOES NOT MEET COVERAGE CRITERIA**.

SPECIMEN VALIDITY TESTING

- 5) Specimen validity testing (e.g., urine specific gravity, urine creatinine, pH, urine oxidant level, genetic identity testing [e.g., NextGen Precision™ Testing]) **DOES NOT MEET COVERAGE CRITERIA**.

NOTES:

Documentation Requirements

The patient's medical record must contain documentation that fully supports the medical necessity for drug testing. This documentation includes, but is not limited to, relevant medical history, physical examination, and results of pertinent diagnostic tests or procedures.

III. Reimbursement

- 1. The following **IS** reimbursed (see complete Coverage Criteria in Letters A and B, Section III above) for:

Reimbursement Policy

- a. Presumptive drug screening based upon appropriate clinical criteria (qualitative, semi-quantitative or quantitative);
 - b. Definitive drug testing (qualitative or quantitative) for up to seven drug classes when the presumptive drug screening meets one of the following criteria:
 - i. The test was negative for prescribed medications, or
 - ii. Positive for a prescription drug with abuse potential which was not prescribed, or
 - iii. Positive for an illegal drug, or
 - iv. A presumptive test does not exist or does not adequately detect the specific drug or metabolite to be tested
 - c. Blood specimens in patients with anuric Chronic Renal Failure.
2. The following **IS NOT REIMBURSED**:
- a. Any AMA definitive drug class codes
 - b. Same-day testing of the same drug or metabolites from two different samples (e.g. both a blood and a urine specimen) by either presumptive or definitive analyses
 - c. Blanket orders or routine standing orders for all patients in the physician's practice
3. Only urine or oral fluid specimens will be covered except blood specimen will be covered for patients with anuric Chronic Renal Failure.
4. Confirmatory/definitive testing should be supported by documentation of rationale in the patient's medical record.
5. More than one presumptive test result per patient per date of service regardless of the number of billing providers **IS NOT REIMBURSED**:
- a. It is not reasonable or necessary for a provider to perform qualitative point-of-care testing and also order presumptive testing from a reference laboratory on the same specimen.
 - b. It is not reasonable or necessary for a provider to perform presumptive immunoassay testing and also order presumptive immunoassay testing from a reference laboratory with or without reflex testing on the same specimen.

IV. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this Policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <https://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

Reimbursement Policy

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

V. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
80305	Drug test(s), presumptive, any number of drug classes, any number of devices or procedures; capable of being read by direct optical observation only (e.g., utilizing immunoassay [eg, dipsticks, cups, cards, or cartridges]), includes sample validation when performed, per date of service
80306	Drug test(s), presumptive, any number of drug classes, any number of devices or procedures; read by instrument assisted direct optical observation (e.g., utilizing immunoassay [eg, dipsticks, cups, cards, or cartridges]), includes sample validation when performed, per date of service
80307	Drug test(s), presumptive, any number of drug classes, any number of devices or procedures; by instrument chemistry analyzers (e.g., utilizing immunoassay [e.g., EIA, ELISA, EMIT, FPIA, IA, KIMS, RIA]), chromatography (e.g., GC, HPLC), and mass spectrometry either with or without chromatography, (e.g., DART, DESI, GC-MS, GC-MS/MS, LC-MS, LC-MS/MS, LDTD, MALDI, TOF) includes sample validation when performed, per date of service
0007U	Drug test(s), presumptive, with definitive confirmation of positive results, any number of drug classes, urine, includes specimen verification including DNA authentication in comparison to buccal DNA, per date of service Proprietary test: ToxProtect Lab/Manufacturer: Genotox Laboratories LTD
0011U	Prescription drug monitoring, evaluation of drugs present by LC-MS/MS, using oral fluid, reported as a comparison to an estimated steady-state range, per date of service including all drug compounds and metabolites Proprietary test: Cordant CORE™ Lab/Manufacturer: Cordant Health Solutions
0051U	Prescription drug monitoring, evaluation of drugs present by LC-MS/MS, urine, 31 drug panel, reported as quantitative results, detected or not detected, per date of service Proprietary test: UCompliDx Lab/Manufacturer: Elite Medical Laboratory Solutions, LLC (LDT)
0054U	Prescription drug monitoring, 14 or more classes of drugs and substances, definitive tandem mass spectrometry with chromatography, capillary blood, quantitative report with therapeutic and toxic ranges, including steady-state

Reimbursement Policy

CPT	Code Description
	range for the prescribed dose when detected, per date of service Proprietary test: AssuranceRx Micro Serum Lab/Manufacturer: Firstox Laboratories, LLC
0079U	Comparative DNA analysis using multiple selected single-nucleotide polymorphisms (SNPs), urine and buccal DNA, for specimen identity verification Proprietary test: ToxLok™ Lab/Manufacturer: InSource Diagnostics
0082U	Drug test(s), definitive, 90 or more drugs or substances, definitive chromatography with mass spectrometry, and presumptive, any number of drug classes, by instrument chemistry analyzer (utilizing immunoassay), urine, report of presence or absence of each drug, drug metabolite or substance with description and severity of significant interactions per date of service Proprietary test: NextGen Precision™ Testing Lab/Manufacturer: Precision Diagnostics LBN Precision Toxicology, LLC
0093U	Prescription drug monitoring, evaluation of 65 common drugs by LC-MS/MS, urine, each drug reported detected or not detected Proprietary test: ComplyRX Lab/Manufacturer: Claro Labs
0227U	Drug assay, presumptive, 30 or more drugs or metabolites, urine, liquid chromatography with tandem mass spectrometry (LC-MS/MS) using multiple reaction monitoring (MRM), with drug or metabolite description, includes sample validation Proprietary Test: Comprehensive Screen Lab/Manufacturer: Aspent Health
0328U	Drug assay, definitive, 120 or more drugs and metabolites, urine, quantitative liquid chromatography with tandem mass spectrometry (LC-MS/MS), includes specimen validity and algorithmic analysis describing drug or metabolite and presence or absence of risks for a significant patient-adverse event, per date of service Proprietary test: CareView360 Lab/Manufacturer: Newstar Medical Laboratories, LLC
G0480	Drug test(s), definitive, utilizing (1) drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (e.g., IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (e.g., alcohol dehydrogenase)), (2) stable isotope or other universally recognized internal standards in all samples (e.g., to control for matrix effects,

Reimbursement Policy

CPT	Code Description
	interferences and variations in signal strength), and (3) method or drug-specific calibration and matrix-matched quality control material (e.g., to control for instrument variations and mass spectral drift); qualitative or quantitative, all sources, includes specimen validity testing, per day; 1-7 drug class(es), including metabolite(s) if performed
G0481	Drug test(s), definitive, utilizing (1) drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (e.g., IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (e.g., alcohol dehydrogenase)), (2) stable isotope or other universally recognized internal standards in all samples (e.g., to control for matrix effects, interferences and variations in signal strength), and (3) method or drug-specific calibration and matrix-matched quality control material (e.g., to control for instrument variations and mass spectral drift); qualitative or quantitative, all sources, includes specimen validity testing, per day; 8-14 drug class(es), including metabolite(s) if performed
G0482	Drug test(s), definitive, utilizing (1) drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (e.g., IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (e.g., alcohol dehydrogenase)), (2) stable isotope or other universally recognized internal standards in all samples (e.g., to control for matrix effects, interferences and variations in signal strength), and (3) method or drug-specific calibration and matrix-matched quality control material (e.g., to control for instrument variations and mass spectral drift); qualitative or quantitative, all sources, includes specimen validity testing, per day; 15-21 drug class(es), including metabolite(s) if performed
G0483	Drug test(s), definitive, utilizing (1) drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including, but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem and excluding immunoassays (e.g., IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (e.g., alcohol dehydrogenase)), (2) stable isotope or other universally recognized internal standards in all samples (e.g., to control for matrix effects, interferences and variations in signal strength), and (3) method or drug-specific calibration and matrix-matched quality control material (e.g., to control for instrument variations and mass spectral drift); qualitative or

Reimbursement Policy

CPT	Code Description
	quantitative, all sources, includes specimen validity testing, per day; 22 or more drug class(es), including metabolite(s) if performed
G0659	Drug test(s), definitive, utilizing drug identification methods able to identify individual drugs and distinguish between structural isomers (but not necessarily stereoisomers), including but not limited to GC/MS (any type, single or tandem) and LC/MS (any type, single or tandem), excluding immunoassays (e.g., IA, EIA, ELISA, EMIT, FPIA) and enzymatic methods (e.g., alcohol dehydrogenase), performed without method or drug-specific calibration, without matrix-matched quality control material, or without use of stable isotope or other universally recognized internal standard(s) for each drug, drug metabolite or drug class per specimen; qualitative or quantitative, all sources, includes specimen validity testing, per day, any number of drug classes

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

VI. Evidence-based Scientific References

- AACC. (2017). *Using Clinical Laboratory Tests to Monitor Drug Therapy in Pain Management Patients*. <https://www.aacc.org/science-and-practice/practice-guidelines/using-clinical-laboratory-tests-to-monitor-drug-therapy-in-pain-management-patients>
- AAFP. (2020). *Clinical Preventive Service Recommendation Opioid Use Disorder (OUD): Screening*. AAFP. Retrieved 1/26 from <https://www.aafp.org/family-physician/patient-care/clinical-recommendations/all-clinical-recommendations/oud.html>
- AAN. (2018). *Diagnostic Assessment of The Child With Status Epilepticus*. <https://www.aan.com/Guidelines/home/GuidelineDetail/234>
- AAPM. (2013). *Use of Opioids for the Treatment of Chronic Pain*. <https://www.ashp.org/-/media/assets/pharmacy-practice/resource-centers/pain-management-toolkit/docs/use-of-opioids-for-the-treatment-of-chronic-pain.ashx>
- AATOD. (2017). *Guidelines for Addressing Benzodiazepine Use in Opioid Treatment Programs (OTPs)*. <https://www.aatod.org/advocacy/policy-statements/guidelines-for-addressing-benzodiazepine-use-in-opioid-treatment-programs-otps-april-6-2017/>
- ACOG. (2017). *Opioid Use and Opioid Use Disorder in Pregnancy*. <https://www.acog.org/-/media/Committee-Opinions/Committee-on-Obstetric-Practice/co711.pdf>
- ALFA. (2019). https://www.cliawaived.com/web/items/pdf/ALF_03_3152_1_Panel_Drug_Test_Insert~493file1.pdf

Reimbursement Policy

- Algren, D. A., & Christian, M. R. (2015). Buyer Beware: Pitfalls in Toxicology Laboratory Testing. *Mo Med*, 112(3), 206-210.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6170116/>
- Altunkaya, D., & Smith, R. N. (1990). Aberrant radioimmunoassay results for cannabinoids in urine. *Forensic Sci Int*, 47(3), 195-205.
- AMDG. (2015). *Interagency Guideline on Prescribing Opioids for Pain*.
<http://www.agencymeddirectors.wa.gov/Files/2015AMDGOpioidGuideline.pdf>
- AMDG. (2018). *Supplemental Guidance on Prescribing Opioids for Postoperative Pain*.
<http://agencymeddirectors.wa.gov/Files/FinalSupBreeAMDGPostopPain091318wcover.pdf>
- APA. (2016). *THE AMERICAN PSYCHIATRIC ASSOCIATION PRACTICE GUIDELINES FOR THE PSYCHIATRIC EVALUATION OF ADULTS*.
<https://psychiatryonline.org/doi/pdf/10.1176/appi.books.9780890426760>
- Argoff, C. E., Alford, D. P., Fudin, J., Adler, J. A., Bair, M. J., Dart, R. C., Gandolfi, R., McCarberg, B. H., Stanos, S. P., Gudim, J. A., Polomano, R. C., & Webster, L. R. (2018). Rational Urine Drug Monitoring in Patients Receiving Opioids for Chronic Pain: Consensus Recommendations. *Pain Med*, 19(1), 97-117. <https://doi.org/10.1093/pm/pnx285>
- ASAM. (2013). *Drug Testing: A White Paper of the American Society of Addiction Medicine (ASAM)*. <https://www.cmm.com.au/files/uploads/resources/20170817102442drug-testing-a-white-paper-by-asam.pdf>
- ASAM. (2017). *Appropriate Use of Drug Testing in Clinical Addiction Medicine*.
file:///C:/Users/AHCS8430/OneDrive%20-%20AVALON%20HEALTH%20SERVICES,%20LLC/Desktop/ASAM%20Guidelines%20Appropriate%20Use%20of%20Drug%20Testing%20in%20Clinical%20Addiction%20Medicine.pdf
- Becker, W., & Starrels, J. (2022, 11/7/2022). *Prescription drug misuse: Epidemiology, prevention, identification, and management*.
<https://www.uptodate.com/contents/prescription-drug-misuse-epidemiology-prevention-identification-and-management>
- Bertron, J. L., Seto, M., & Lindsley, C. W. (2018). DARK Classics in Chemical Neuroscience: Phencyclidine (PCP). *ACS Chem Neurosci*, 9(10), 2459-2474.
<https://doi.org/10.1021/acschemneuro.8b00266>
- Blank, A., Hellstern, V., Schuster, D., Hartmann, M., Matthee, A. K., Burhenne, J., Haefeli, W. E., & Mikus, G. (2009). Efavirenz treatment and false-positive results in benzodiazepine screening tests. *Clin Infect Dis*, 48(12), 1787-1789. <https://doi.org/10.1086/599109>
- Böttcher, M., Lierheimer, S., Peschel, A., & Beck, O. (2019). Detection of heroin intake in patients in substitution treatment using oral fluid as specimen for drug testing. *Drug Alcohol Depend*, 198, 136-139. <https://doi.org/10.1016/j.drugalcdep.2019.02.011>
- Brahm, N. C., Yeager, L. L., Fox, M. D., Farmer, K. C., & Palmer, T. A. (2010). Commonly prescribed medications and potential false-positive urine drug screens. *Am J Health Syst Pharm*, 67(16), 1344-1350. <https://doi.org/10.2146/ajhp090477>
- CDC. (2015, June 2015). *Nationwide Trends*.
<https://www.drugabuse.gov/publications/drugfacts/nationwide-trends>

Reimbursement Policy

- CDC. (2017, 02/09/2017). *Opioid Data Analysis and Resources*. Centers for Disease Control and Prevention. Retrieved 10/01/2018 from <https://www.cdc.gov/drugoverdose/data/analysis.html>
- CDC. (2018a, December 2018). *Monitoring the Future Survey: High School and Youth Trends*. CDC. <https://www.drugabuse.gov/publications/drugfacts/monitoring-future-survey-high-school-youth-trends>
- CDC. (2018b). *Quality Improvement and Care Coordination: Implementing the CDC Guideline for Prescribing Opioids for Chronic Pain* Retrieved 2/1/2021 from <https://www.cdc.gov/drugoverdose/pdf/prescribing/CDC-DUIP-QualityImprovementAndCareCoordination-508.pdf>
- CDC. (2019). *Other Drugs*. <https://www.cdc.gov/drugoverdose/data/otherdrugs.html>
- CDC. (2022a). *Drug Overdose Deaths*. <https://www.cdc.gov/drugoverdose/deaths/index.html>
- CDC. (2022b). *U.S. Opioid Dispensing Rate Maps*. <https://www.cdc.gov/drugoverdose/rxrate-maps/index.html>
- Chou, R., Fanciullo, G. J., Fine, P. G., Adler, J. A., Ballantyne, J. C., Davies, P., Donovan, M. I., Fishbain, D. A., Foley, K. M., Fudin, J., Gilson, A. M., Kelter, A., Mauskop, A., O'Connor, P. G., Passik, S. D., Pasternak, G. W., Portenoy, R. K., Rich, B. A., Roberts, R. G., . . . Miaskowski, C. (2009). Clinical Guidelines for the Use of Chronic Opioid Therapy in Chronic Noncancer Pain. *The Journal of Pain*, 10(2), 113-130.e122. <https://doi.org/10.1016/j.jpain.2008.10.008>
- Christo, P. J., Manchikanti, L., Ruan, X., Bottros, M., Hansen, H., Solanki, D. R., Jordan, A. E., & Colson, J. (2011). Urine drug testing in chronic pain. *Pain Physician*, 14(2), 123-143. <https://www.painphysicianjournal.com/current/pdf?article=MTQ0NA%3D%3D&journal=60>
- Chua, I., Petrides, A. K., Schiff, G. D., Ransohoff, J. R., Kantartjis, M., Streid, J., Demetriou, C. A., & Melanson, S. E. F. (2020). Provider Misinterpretation, Documentation, and Follow-Up of Definitive Urine Drug Testing Results. *J Gen Intern Med*, 35(1), 283-290. <https://doi.org/10.1007/s11606-019-05514-5>
- Cone, E. J., Bigelow, G. E., Herrmann, E. S., Mitchell, J. M., LoDico, C., Flegel, R., & Vandrey, R. (2015). Non-smoker exposure to secondhand cannabis smoke. I. Urine screening and confirmation results. *J Anal Toxicol*, 39(1), 1-12. <https://doi.org/10.1093/jat/bku116>
- Couto, J. E., Webster, L., Romney, M. C., Leider, H. L., & Linden, A. (2009). Use of an algorithm applied to urine drug screening to assess adherence to an oxycontin regimen. *J Opioid Manag*, 5(6), 359-364.
- Couto, J. E., Webster, L., Romney, M. C., Leider, H. L., & Linden, A. (2011). Use of an algorithm applied to urine drug screening to assess adherence to a hydrocodone regimen. *J Clin Pharm Ther*, 36(2), 200-207. <https://doi.org/10.1111/j.1365-2710.2010.01236.x>
- DOD. (2017). *VA/DoD Clinical Practice Guideline for Opioid Therapy For Chronic Pain*. <https://www.healthquality.va.gov/guidelines/Pain/cot/VADoDOTCPG022717.pdf>
- DOD. (2018). *VA/DOD Clinical Practice Guideline for The Management Of Pregnancy*. <https://www.healthquality.va.gov/guidelines/WH/up/VADoDPregnancyCPG4102018.pdf>
- Dowell, D., Haegerich, T. M., & Chou, R. (2016). CDC Guideline for Prescribing Opioids for Chronic Pain - United States, 2016. *MMWR Recomm Rep*, 65(1), 1-49. <https://doi.org/10.15585/mmwr.rr6501e1>

Reimbursement Policy

- Drake, L. R., & Scott, P. J. H. (2018). DARK Classics in Chemical Neuroscience: Cocaine. *ACS Chem Neurosci*, 9(10), 2358-2372. <https://doi.org/10.1021/acscchemneuro.8b00117>
- DVA, & DOD. (2015). *VA/DoD Clinical Practice Guideline for The Management Of Substance Use Disorders*.
<https://www.healthquality.va.gov/guidelines/MH/sud/VADoDSUDCPGRevised22216.pdf>
- DWD. (2013). *Chronic Opioid Clinical Management Guidelines for Wisconsin Worker's Compensation Patient Care*. Department of Workforce Development, State of Wisconsin.
<https://dwd.wisconsin.gov/wc/medical/pdf/CHRONIC%20OPIOID%20CLINICAL%20MANAGEMENT%20GUIDELINES%20.pdf>
- Eaton, K., & Lyman, G. (2022, 10/21/2022). *Dosing of anticancer agents in adults*.
<https://www.uptodate.com/contents/dosing-of-anticancer-agents-in-adults>
- Eskridge, K. D., & Guthrie, S. K. (1997). Clinical issues associated with urine testing of substances of abuse. *Pharmacotherapy*, 17(3), 497-510.
- Farkas, A., Lipanot, K., & Sherman, K. (2021). Routine Laboratory Screening for Acetaminophen and Salicylate Ingestion in Preadmission Psychiatric Patients Is Unnecessary. *Ann Emerg Med*, 77(6), 604-612.
<https://doi.org/10.1016/j.annemergmed.2021.01.027>
- FDA. (2018). *DRI Cocaine Metabolite Assay*.
https://www.accessdata.fda.gov/cdrh_docs/pdf18/K181499.pdf
- FDA. (2019). *510(k) Substantial Equivalence Determination Decision Summary Assay Only Template* https://www.accessdata.fda.gov/cdrh_docs/reviews/K112395.pdf
- FSMB. (2017). *Guidelines for the Chronic Use of Opioid Analgesics*. Federation of State Medical Boards. Retrieved 10/01/2018 from
https://www.fsmb.org/globalassets/advocacy/policies/opioid_guidelines_as_adopted_april-2017_final.pdf
- Fucci, N. (2012). False positive results for amphetamine in urine of a patient with diabetes mellitus. *Forensic Sci Int*, 223(1-3), e60. <https://doi.org/10.1016/j.forsciint.2012.08.010>
- Grant, C. N., & Bélanger, R. E. (2017). Position Statement: Cannabis and Canada's children and youth. *Pediatric Child Health*, 22(2), 98-102.
<https://www.cps.ca/en/documents/position/cannabis-children-and-youth>
- Gray, C., Korczak, D., Andrews, D., & Bélanger, S. A. (2018). ADHD in children and youth: Part 1—Etiology, diagnosis, and comorbidity. *Paediatrics & Child Health*, 23(7), 447-453.
<https://doi.org/10.1093/pch/pxy109>
- Greller, H., & Gupta, A. (2022, 07/11/2022). *Benzodiazepine poisoning and withdrawal*.
<https://www.uptodate.com/contents/benzodiazepine-poisoning-and-withdrawal>
- Herrmann, E. S., Cone, E. J., Mitchell, J. M., Bigelow, G. E., LoDico, C., Flegel, R., & Vandrey, R. (2015). Non-smoker exposure to secondhand cannabis smoke II: Effect of room ventilation on the physiological, subjective, and behavioral/cognitive effects. *Drug Alcohol Depend*, 151, 194-202. <https://doi.org/10.1016/j.drugalcdep.2015.03.019>
- HHS. (2017). *New HHS Guidelines for Drug Testing Panels Effective 10/1/17*.
<https://www.federalregister.gov/documents/2017/01/23/2017-00979/mandatory-guidelines-for-federal-workplace-drug-testing-programs#h-114>

Reimbursement Policy

- Hoffman, R. (2022, 9/21/2022). *Testing for drugs of abuse (DOAs)*. Wolters Kluwer.
<https://www.uptodate.com/contents/testing-for-drugs-of-abuse-doas>
- Jamison, R. N., Ross, E. L., Michna, E., Chen, L. Q., Holcomb, C., & Wasan, A. D. (2010). Substance misuse treatment for high-risk chronic pain patients on opioid therapy: a randomized trial. *Pain*, 150(3), 390-400. <https://doi.org/10.1016/j.pain.2010.02.033>
- Jannetto, P., Bratanow, N., Clark, W., Hamill-Ruth, R., Hammett-Stabler, C., Huestis, M., Kassed, C., McMillin, G., Melanson, S., & Langman, L. (2017). *Using Clinical Laboratory Tests to Monitor Drug Therapy in Pain Management Patients*. AACC. Retrieved 10/01/2018 from <https://www.aacc.org/science-and-practice/practice-guidelines/using-clinical-laboratory-tests-to-monitor-drug-therapy-in-pain-management-patients>
- Jannetto, P. J., & Langman, L. J. (2018). Using Clinical Laboratory Tests to Monitor Drug Therapy in Pain Management Patients. *The Journal of Applied Laboratory Medicine: An AACC Publication*, 2(4), 471-472. <https://doi.org/10.1373/jalm.2017.025304>
- Jarvis, M., Williams, J., Hurford, M., Lindsay, D., Lincoln, P., Giles, L., Luongo, P., & Safarian, T. (2017). Appropriate Use of Drug Testing in Clinical Addiction Medicine. *J Addict Med*, 11(3), 163-173. <https://doi.org/10.1097/adm.0000000000000323>
- Jones, C. M., Mack, K. A., & Paulozzi, L. J. (2013). Pharmaceutical overdose deaths, United States, 2010. *Jama*, 309(7), 657-659. <https://doi.org/10.1001/jama.2013.272>
- Jones, J. (2016, 07/28/2016). *Clinical vs. Forensic: The Differences Cost More Than Just Money*. United States Drug Testing Laboratories. Retrieved 02/05/2019 from <http://www.usdtl.com/media/mediaarticles/clinical-vs-forensic-the-differences-cost-more-than-just-money>
- Kale, N. (2019). Urine Drug Tests: Ordering and Interpreting Results. *Am Fam Physician*, 99(1), 33-39. <https://www.aafp.org/afp/2019/0101/p33.pdf>
- Kampman, K. (2022, 7/11/2022). *Approach to treatment of stimulant use disorder in adults*. <https://www.uptodate.com/contents/approach-to-treatment-of-stimulant-use-disorder-in-adults>
- Kandel, D. B., Hu, M. C., Griesler, P., & Wall, M. (2017). Increases from 2002 to 2015 in prescription opioid overdose deaths in combination with other substances. *Drug Alcohol Depend*, 178, 501-511. <https://doi.org/10.1016/j.drugalcdep.2017.05.047>
- Katz, N. P., Sherburne, S., Beach, M., Rose, R. J., Vielguth, J., Bradley, J., & Fanciullo, G. J. (2003). Behavioral monitoring and urine toxicology testing in patients receiving long-term opioid therapy. *Anesth Analg*, 97(4), 1097-1102, table of contents.
- Katzman, M. A., Bleau, P., Blier, P., Chokka, P., Kjernisted, K., Van Ameringen, M., Canadian Anxiety Guidelines Initiative Group on behalf of the Anxiety Disorders Association of Canada/Association Canadienne des troubles, a., McGill, U., Antony, M. M., Bouchard, S., Brunet, A., Flament, M., Grigoriadis, S., Mendlowitz, S., O'Connor, K., Rabheru, K., Richter, P. M. A., Robichaud, M., & Walker, J. R. (2014). Canadian clinical practice guidelines for the management of anxiety, posttraumatic stress and obsessive-compulsive disorders. *BMC psychiatry*, 14 Suppl 1(Suppl 1), S1-S1. <https://doi.org/10.1186/1471-244X-14-S1-S1>

Reimbursement Policy

- Kell, M. J. (1994). Utilization of plasma and urine methadone concentrations to optimize treatment in maintenance clinics: I. Measurement techniques for a clinical setting. *J Addict Dis*, 13(1), 5-26. https://doi.org/10.1300/J069v13n01_02
- Knezevic, N. N., Khan, O. M., Beiranvand, A., & Candido, K. D. (2017). Repeated Quantitative Urine Toxicology Analysis May Improve Chronic Pain Patient Compliance with Opioid Therapy. *Pain Physician*, 20(2s), S135-s145.
- Krasowski, M. D., McMillin, G. A., Melanson, S. E. F., Dizon, A., Magnani, B., & Snozek, C. L. H. (2020). Interpretation and Utility of Drug of Abuse Screening Immunoassays: Insights From Laboratory Drug Testing Proficiency Surveys. *Arch Pathol Lab Med*, 144(2), 177-184. <https://doi.org/10.5858/arpa.2018-0562-CP>
- Langman, L. J., & Jannetto, P. J. (2018). *Using Clinical Laboratory Tests to Monitor Drug Therapy in Pain Management Patients*. <https://www.aacc.org/-/media/Files/Science-and-Practice/Practice-Guidelines/Pain-Management/LMPGPain-Management20171220.pdf>
- Levine, B. S., & Smith, M. L. (1990). Effects of diphenhydramine on immunoassays of phencyclidine in urine. *Clin Chem*, 36(6), 1258.
- Ly, B. T., Thornton, S. L., Buono, C., Stone, J. A., & Wu, A. H. (2012). False-positive urine phencyclidine immunoassay screen result caused by interference by tramadol and its metabolites. *Ann Emerg Med*, 59(6), 545-547. <https://doi.org/10.1016/j.annemergmed.2011.08.013>
- Manchikanti, L., Abdi, S., Atluri, S., Balog, C. C., Benyamin, R. M., Boswell, M. V., Brown, K. R., Bruel, B. M., Bryce, D. A., Burks, P. A., Burton, A. W., Calodney, A. K., Caraway, D. L., Cash, K. A., Christo, P. J., Damron, K. S., Datta, S., Deer, T. R., Diwan, S., . . . Wargo, B. W. (2012). American Society of Interventional Pain Physicians (ASIPP) guidelines for responsible opioid prescribing in chronic non-cancer pain: Part 2--guidance. *Pain Physician*, 15(3 Suppl), S67-116.
- Manchikanti, L., Kaye, A. M., Knezevic, N. N., McAnally, H., Slavin, K., Trescot, A. M., Blank, S., Pampati, V., Abdi, S., Grider, J. S., Kaye, A. D., Manchikanti, K. N., Corder, H., Gharibo, C. G., Harned, M. E., Albers, S. L., Atluri, S., Aydin, S. M., Bakshi, S., . . . Hirsch, J. A. (2017). Responsible, Safe, and Effective Prescription of Opioids for Chronic Non-Cancer Pain: American Society of Interventional Pain Physicians (ASIPP) Guidelines. *Pain Physician*, 20(2s), S3-s92.
- Manchikanti, L., Malla, Y., Wargo, B. W., Cash, K. A., Pampati, V., Damron, K. S., McManus, C. D., & Brandon, D. E. (2010). Protocol for accuracy of point of care (POC) or in-office urine drug testing (immunoassay) in chronic pain patients: a prospective analysis of immunoassay and liquid chromatography tandem mass spectrometry (LC/MS/MS). *Pain Physician*, 13(1), E1-e22. <https://www.painphysicianjournal.com/current/pdf?article=MTMwMg%3D%3D&journal=53>
- McClellan, J., & Stock, S. (2013). Practice Parameter for the Assessment and Treatment of Children and Adolescents With Schizophrenia. *Journal of the American Academy of Child & Adolescent Psychiatry*, 52(9), 976-990. <https://doi.org/10.1016/j.jaac.2013.02.008>
- McEvoy, J., Millet, R. A., Dretchen, K., Morris, A. A., Corwin, M. J., & Buckley, P. (2014). Quantitative levels of aripiprazole parent drug and metabolites in urine.

Reimbursement Policy

- Psychopharmacology (Berl)*, 231(23), 4421-4428. <https://doi.org/10.1007/s00213-014-3781-1>
- Michna, E., Jamison, R. N., Pham, L. D., Ross, E. L., Janfaza, D., Nedeljkovic, S. S., Narang, S., Palombi, D., & Wasan, A. D. (2007). Urine toxicology screening among chronic pain patients on opioid therapy: frequency and predictability of abnormal findings. *Clin J Pain*, 23(2), 173-179. <https://doi.org/10.1097/AJP.0b013e31802b4f95>
- Microgenics. (2015). *CEDIA(r) Phencyclidine (PCP) Assay*. Microgenics Corporation. <http://tools.thermofisher.com/content/sfs/manuals/10007400-CEDIA-Phencyclidine-PCP-Assay-EN.pdf>
- Microgenics. (2016). *DRI(r) Amphetamines Assay*. Microgenics Corporation. Retrieved 02/12/2019 from <http://tools.thermofisher.com/content/sfs/manuals/0138-DRI-Amphetamines-Assay-EN.pdf>
- Moeller, K. E., Kissack, J. C., Atayee, R. S., & Lee, K. C. (2017). Clinical Interpretation of Urine Drug Tests: What Clinicians Need to Know About Urine Drug Screens. *Mayo Clin Proc*, 92(5), 774-796. <https://doi.org/10.1016/j.mayocp.2016.12.007>
- Nafziger, A. N., & Bertino, J. S., Jr. (2009). Utility and application of urine drug testing in chronic pain management with opioids. *Clin J Pain*, 25(1), 73-79. <https://doi.org/10.1097/AJP.0b013e31817e13cc>
- National Center for Drug Abuse Statistics. (2022). *Drug Abuse Statistics*. <https://drugabusestatistics.org/>
- Nelson, L., & Odujebi, O. (2022, 6/7/2022). *Cocaine: Acute intoxication*. <https://www.uptodate.com/contents/cocaine-acute-intoxication>
- NICE. (2018, April 2018). *Epilepsies: diagnosis and management*. <https://www.nice.org.uk/guidance/cg137/chapter/1-Guidance>
- NICE. (2020). *Epilepsies: diagnosis and management*. <https://www.nice.org.uk/guidance/cg137/chapter/1-Guidance>
- Nuckols, T. K., Anderson, L., Popescu, I., Diamant, A. L., Doyle, B., Di Capua, P., & Chou, R. (2014). Opioid prescribing: a systematic review and critical appraisal of guidelines for chronic pain. *Ann Intern Med*, 160(1), 38-47. <https://doi.org/10.7326/0003-4819-160-1-201401070-00732>
- OASAS. (2021). *Guidance on Toxicology Use in OASAS Certified Programs*. https://oasas.ny.gov/system/files/documents/2021/11/oasas-toxicology-guidance_0.pdf
- Ordean, A., Wong, S., & Graves, L. (2017). No. 349-Substance Use in Pregnancy. *J Obstet Gynaecol Can*, 39(10), 922-937.e922. <https://doi.org/10.1016/j.jogc.2017.04.028>
- Owen, G. T., Burton, A. W., Schade, C. M., & Passik, S. (2012). Urine drug testing: current recommendations and best practices. *Pain Physician*, 15(3 Suppl), E119-133.
- Owusu Obeng, A., Hamadeh, I., & Smith, M. (2017). Review of Opioid Pharmacogenetics and Considerations for Pain Management. *Pharmacotherapy*, 37(9), 1105-1121. <https://doi.org/10.1002/phar.1986>
- Palamar, J. J., Le, A., Guarino, H., & Mateu-Gelabert, P. (2019). A comparison of the utility of urine- and hair testing in detecting self-reported drug use among young adult opioid users. *Drug Alcohol Depend*, 200, 161-167. <https://doi.org/10.1016/j.drugalcdep.2019.04.008>

Reimbursement Policy

- Pesce, A., Krock, K., Ritz, D., Cua, A., Thomas, R., & Nickley, J. (2018). Observations on 6-MAM (6-Monoacetylmorphine) in Urine. *J Clin Toxicol*, 8(393), 2161-0495.1000393.
- Pesce, A., West, C., Egan City, K., & Strickland, J. (2012). Interpretation of urine drug testing in pain patients. *Pain Med*, 13(7), 868-885. <https://doi.org/10.1111/j.1526-4637.2012.01350.x>
- Phan, H. M., Yoshizuka, K., Murry, D. J., & Perry, P. J. (2012). Drug testing in the workplace. *Pharmacotherapy*, 32(7), 649-656. <https://doi.org/10.1002/j.1875-9114.2011.01089.x>
- PMFT. (2019). *Pain Management Best Practices Inter-Agency Task Force Report: Updates, Gaps, Inconsistencies, and Recommendations- Final Report*
<https://www.hhs.gov/sites/default/files/pmtf-final-report-2019-05-23.pdf>
- Rengarajan, A., & Mullins, M. E. (2013). How often do false-positive phencyclidine urine screens occur with use of common medications? *Clin Toxicol (Phila)*, 51(6), 493-496.
<https://doi.org/10.3109/15563650.2013.801982>
- Rollins, D. E., Jennison, T. A., & Jones, G. (1990). Investigation of interference by nonsteroidal anti-inflammatory drugs in urine tests for abused drugs. *Clin Chem*, 36(4), 602-606.
- SAMHSA. (2007). *Guidelines for the Accreditation of Opioid Treatment Programs*
https://www.samhsa.gov/sites/default/files/programs_campaigns/medication_assisted/2007-otp-accreditation-guidelines.pdf
- SAMHSA. (2015). *FEDERAL GUIDELINES FOR OPIOID TREATMENT PROGRAMS*.
<https://store.samhsa.gov/system/files/pep15-fedguideotp.pdf>
- SAMHSA. (2017). *Key Substance Use and Mental Health Indicators in the United States: Results from the 2016 National Survey on Drug Use and Health*.
<https://www.samhsa.gov/data/sites/default/files/NSDUH-FFR1-2016/NSDUH-FFR1-2016.pdf>
- SAMHSA. (2020). *Mandatory Guidelines for Federal Workplace Drug Testing Programs—Oral/Fluid*
https://www.samhsa.gov/sites/default/files/programs_campaigns/division_workplace_programs/final-mg-oral-fluid.pdf
- SAMHSA. (2022). *Mandatory Guidelines for Federal Workplace Drug Testing Programs*. Federal Register. <https://www.federalregister.gov/documents/2022/04/07/2022-06884/mandatory-guidelines-for-federal-workplace-drug-testing-programs#sectno-reference-2.2>
- Smith, P. E., & McBride, A. (1999). Illicit drugs and seizures. *Seizure*, 8(8), 441-443.
<https://doi.org/10.1053/seiz.1999.0346>
- Snyder, M. L., Fantz, C. R., & Melanson, S. (2017). Immunoassay-Based Drug Tests Are Inadequately Sensitive for Medication Compliance Monitoring in Patients Treated for Chronic Pain. *Pain Physician*, 20(2s), Se1-se9.
<https://www.painphysicianjournal.com/current/pdf?article=NDIwNw%3D%3D&journal=103>
- Starrels, J. L., Becker, W. C., Alford, D. P., Kapoor, A., Williams, A. R., & Turner, B. J. (2010). Systematic review: treatment agreements and urine drug testing to reduce opioid misuse in patients with chronic pain. *Ann Intern Med*, 152(11), 712-720. <https://doi.org/10.7326/0003-4819-152-11-201006010-00004>

Reimbursement Policy

- Trescot, A. M., Datta, S., Lee, M., & Hansen, H. (2008). Opioid pharmacology. *Pain Physician*, 11(2 Suppl), S133-153.
- Vohra, V., Marraffa, J. M., Wojcik, S. M., & Eggleston, W. (2019). An assessment of urine THC immunoassay in healthy volunteers receiving an oral proton-pump inhibitor. *Clin Toxicol (Phila)*, 1-3. <https://doi.org/10.1080/15563650.2019.1662917>
- Vopat, M. L., Messamore, W. G., Trent, J. J., Schmanke, K. E., Zackula, R., Yang, S. Y., & Bhargava, T. (2020). Urine Screening for Opioid and Illicit Drugs in the Total Joint Arthroplasty Population. *Kans J Med*, 13, 71-76.
- Weaver, M. F. (2015). Prescription Sedative Misuse and Abuse. *Yale J Biol Med*, 88(3), 247-256. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4553644/>
- WFSBP. (2015). *World Federation of Societies of Biological Psychiatry (WFSBP) Guidelines for Biological Treatment of Schizophrenia Part 3: Update 2015 Management of special circumstances: Depression, Suicidality, substance use disorders and pregnancy and lactation* http://www.wfsbp.org/fileadmin/user_upload/Treatment_Guidelines/Hasan_et_al_2015_.pdf
- WHO. (2016). *mhGAP Intervention Guide*. <https://apps.who.int/iris/bitstream/handle/10665/250239/9789241549790-eng.pdf?sequence=1>
- Wilfong, A. (2021, 11/23/2021). *Management of convulsive status epilepticus in children*. <https://www.uptodate.com/contents/management-of-convulsive-status-epilepticus-in-children>
- Wondfo. (2020). *Drug Tests Strip*. <http://wondfousa.com/test/strip/>
- Wong, S., Ordean, A., & Kahan, M. (2011). Substance use in pregnancy. *J Obstet Gynaecol Can*, 33(4), 367-384. [https://doi.org/10.1016/s1701-2163\(16\)34855-1](https://doi.org/10.1016/s1701-2163(16)34855-1)
- Xlar. (2002). *THC One Step Marijuana Test Strip Package Insert*. xlar.com/I/THC-STRIP-2.pdf

VII. Revision History

Revision Date	Summary of Changes
03/06/2024	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria.
05/31/2023	Off-Cycle Review: Proprietary test from Newstar Medical (RiskViewRx) that was called out in CC4 was replaced by CareView360. No further updates outside of the CC. Removed CPT code 0143U – 0150U (deletion effective 7/1/2023)
03/01/2023	Reviewed and Updated: Updated the background, guidelines and recommendations, and evidence-based scientific references. Literature review did not necessitate any modifications to coverage criteria. The following edits were made for clarity: All CC edited for clarity and consistency. Removed CC3 “3) Confirmatory/definitive qualitative or quantitative drug testing DOES NOT MEET COVERAGE CRITERIA when laboratory-based definitive

Reimbursement Policy

	drug testing is requested without any prior presumptive screening test results indicating the clinical utility to confirm those results.” Added CPT 0328U.
--	---