

Reimbursement Policy

Human Immunodeficiency Virus (HIV)

[POLICY DESCRIPTION](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#)

I. Policy Description

Human immunodeficiency virus (HIV) is an RNA retrovirus that infects human immune cells, specifically CD4 cells, causing progressive deterioration of the immune system ultimately leading to acquired immune deficiency syndrome (AIDS) characterized by susceptibility to opportunistic infections and HIV-related cancers.¹ HIV-1 is the dominant subtype of HIV infection, but another subtype, HIV-2, is a crucial subtype in certain areas of the world, such as Western Africa.² Terms such as male and female are used when necessary to refer to sex assigned at birth.

II. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the "Applicable State and Federal Regulations" section of this policy document.

- 1) For individuals 11 to 65 years of age, initial screening for HIV infection with an antigen/antibody combination assay **MEETS COVERAGE CRITERIA**.
- 2) For individuals 11 to 65 years of age, repeat antigen/antibody screening for HIV infection (no more than one test every 90 days) **MEETS COVERAGE CRITERIA**.
- 3) Nucleic acid testing (qualitative or quantitative) for HIV-1 and HIV-2 (no more than one test every month) **MEETS COVERAGE CRITERIA** in **any** of the following situations:
 - a) For individuals for whom initial screening was positive for HIV infection.
 - b) For individuals for whom initial screening was indeterminate for HIV infection.
 - c) For individuals for whom recent exposure is suspected or reported.
- 4) HIV genotyping or phenotyping **MEETS COVERAGE CRITERIA** for **any** of the following situations:
 - a) Prior to initiating doravirine therapy (genotyping and phenotyping is required).

Reimbursement Policy

- b) For individuals who have failed a course of antiviral therapy.
- c) For individuals who have suboptimal viral load reduction.
- d) For individuals who have been noncompliant with therapy.
- e) To guide treatment decisions in individuals with acute or recent infection (within the last 6 months).
- f) For antiretroviral naïve individuals entering treatment.
- g) For all HIV-infected pregnant individuals in the following situations:
 - i) Before initiation of antiretroviral therapy.
 - ii) For those with detectable HIV RNA levels.
- 5) For treatment-experienced individuals on failing regimens who are thought to have multidrug resistance, HIV phenotyping **MEETS COVERAGE CRITERIA**.

Plasma quantification of HIV-1 RNA or HIV-2 RNA (see Note 1) (no more than one test every month) **MEETS COVERAGE CRITERIA** for **any** of the following situations:

- a) For monitoring disease progression in HIV-infected individuals.
- b) For monitoring response to antiretroviral therapy.
- c) For infants younger than 18 months born to HIV-positive mothers (antibody tests may be confounded by maternal antibodies in this time frame).
- d) For predicting maternal-fetal transmission of HIV-1 or HIV-2.

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.

- 6) Routine use of combined genotyping and phenotyping **DOES NOT MEET COVERAGE CRITERIA**.
- 7) Drug susceptibility phenotype prediction using genotypic comparison to known genotypic/phenotypic database **DOES NOT MEET COVERAGE CRITERIA**.

Reimbursement Policy

NOTES:

Note 1: Because differences in absolute HIV copy number are known to occur using different assays, plasma HIV RNA levels should be measured by the same analytical method. A change in assay method may necessitate re-establishment of a baseline.

III. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this Policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <http://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, please visit the applicable state Medicaid website.

Food and Drug Administration (FDA)

The primary RT-PCR tests for HIV-1 have been approved by the FDA:

In May 2007, the FDA approved the Abbott RealTime HIV-1 Amplification Reagent Kit. From the FDA website: “The Abbott RealTime HIV-1 assay is an in vitro reverse transcription-polymerase chain reaction (RT-PCR) assay for the quantitation of HIV-1 on the automated m2000 System in human plasma from HIV-1 infected individuals over the range of 40 to 10,000,000 copies/mL.”⁵²

On May 11, 2007, the FDA approved the COBAS® AmpliPrep/COBAS® TaqMan® HIV-1 Test. From the FDA website: “The COBAS AmpliPrep/COBAS TaqMan HIV-1 is an in vitro nucleic acid amplification test for the quantitation of human immunodeficiency virus (HIV-1) nucleic acid in human plasma, using the COBAS AmpliPrep Instrument for automated sample preparation and the COBAS TaqMan Analyzer or COBAS TaqMan 48 Analyzer for automated amplification and detection. This test is intended for use in conjunction with clinical presentation and other laboratory markers of disease progress for the clinical management of HIV-1 infected patients.”⁵³

In 2016, the FDA approved the Aptima® HIV-1 Quant Assay. From the FDA website: “The Aptima HIV-1 Quant assay is an in vitro nucleic acid amplification test (NAAT) for the quantitation of HIV-1 RNA in human plasma from HIV-1 infected individuals on the fully automated Panther® system. The Aptima HIV-1 Quant assay quantitates HIV-1 RNA groups M, N, and O over the range of 30 to 10,000,000 copies/ mL.” On November 20, 2020, this assay was given an FDA approval for dual use for diagnosis and viral load monitoring for HIV-1.^{4,5}

The following screening antibody tests are FDA-approved to differentiate HIV-1 from HIV-2:

Reimbursement Policy

On August 26, 2019, the FDA approved the Geenius HIV-1/2 Supplemental Assay. From the FDA website: “The Geenius™ HIV 1/2 Supplemental Assay is a single-use immunochromatographic assay for the confirmation and differentiation of individual antibodies to human immunodeficiency virus Types 1 and 2 (HIV-1 and HIV-2) in serum or plasma samples (EDTA, lithium heparin, sodium citrate, and CPD) from blood donors. The Geenius™ HIV 1/2 Supplemental Assay is intended for use as an additional, more specific test for human serum and plasma samples with repeatedly reactive results by an FDA licensed blood donor screening test for antibodies to HIV-1/HIV-2. The results of the Geenius™ HIV 1/2 Supplemental Assay are read and interpreted only with the Geenius™ Reader with dedicated software.” There were 200 known HIV-2 positive samples classified by Geenius, with 77 interpreted as only HIV-2 positive, 108 with HIV-2 with HIV-1 cross reactivity, 12 as undifferentiated, and 3 as HIV-2 indeterminate.⁵⁴

On July 23, 2015, the FDA approved the BioPlex 2200 HIV Ag-Ab assay. From the FDA website: “The BioPlex 2200 HIV Ag-Ab assay is a multiplex flow immunoassay intended for the simultaneous qualitative detection and differentiation of the individual analytes HIV-1 p24 antigen, HIV-1 (groups M and O) antibodies, and HIV-2 antibodies in human serum or plasma (fresh or frozen K2 EDTA, K3 EDTA, lithium heparin, sodium heparin; fresh citrate). This assay is intended as an aid in the diagnosis of infection with HIV-1 and/or HIV-2, including acute (primary) HIV-1 infection. The assay may also be used as an aid in the diagnosis of infection with HIV-1 and/or HIV-2 in pediatric subjects as young as two years of age, and pregnant women.” The test was found to differentiate all 1363 HIV-1 samples correctly and 188 of 200 HIV-2 samples correctly (with 12 “undifferentiated”).⁵⁵

In 2020 and 2022, the FDA approved the Alinity m HIV-1 assay as an in vitro reverse transcription-polymerase chain reaction (RT-PCR) assay for the detection and quantification of HIV-1. It is to be used both for confirmation of HIV-1 infection and for monitoring of HIV-1 infected individuals. From the FDA website: “The Alinity m HIV-1 assay is intended for use to monitor disease prognosis by measuring baseline plasma HIV-1 RNA level and to assess response to antiretroviral treatment by measuring changes in plasma HIV-1 RNA levels. Performance for quantitative monitoring is not established with serum specimens.” The assay can also be used as a supplemental test to confirm HIV-1 in individuals who have “reactive results” with HIV immunoassays.⁵⁶

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

IV. Applicable CPT/HCPCS Procedure Codes

Reimbursement Policy

CPT	Code Description
86689	Antibody; HTLV or HIV antibody, confirmatory test (eg, Western Blot)
86701	Antibody; HIV-1
86702	Antibody; HIV-2
86703	Antibody; HIV-1 and HIV-2, single result
87389	Infectious agent antigen detection by immunoassay technique, (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]) qualitative or semiquantitative; HIV-1 antigen(s), with HIV-1 and HIV-2 antibodies, single result
87390	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; HIV-1
87391	Infectious agent antigen detection by immunoassay technique (eg, enzyme immunoassay [EIA], enzyme-linked immunosorbent assay [ELISA], fluorescence immunoassay [FIA], immunochemiluminometric assay [IMCA]), qualitative or semiquantitative; HIV-2
87534	Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, direct probe technique
87535	Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, amplified probe technique, includes reverse transcription when performed
87536	Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, quantification, includes reverse transcription when performed
87537	Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, direct probe technique
87538	Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, amplified probe technique, includes reverse transcription when performed
87539	Infectious agent detection by nucleic acid (DNA or RNA); HIV-2, quantification, includes reverse transcription when performed

Reimbursement Policy

CPT	Code Description
87806	Infectious agent antigen detection by immunoassay with direct optical (ie, visual) observation; HIV-1 antigen(s), with HIV-1 and HIV-2 antibodies
87900	Infectious agent drug susceptibility phenotype prediction using regularly updated genotypic bioinformatics
87901	Infectious agent genotype analysis by nucleic acid (DNA or RNA); HIV-1, reverse transcriptase and protease regions
87903	Infectious agent phenotype analysis by nucleic acid (DNA or RNA) with drug resistance tissue culture analysis, HIV 1; first through 10 drugs tested
87904	Infectious agent phenotype analysis by nucleic acid (DNA or RNA) with drug resistance tissue culture analysis, HIV 1; each additional drug tested (List separately in addition to code for primary procedure)
87906	Infectious agent genotype analysis by nucleic acid (DNA or RNA); HIV-1, other region (eg, integrase, fusion)
0219U	Infectious agent (human immunodeficiency virus), targeted viral next-generation sequence analysis (ie, protease [PR], reverse transcriptase [RT], integrase [INT]), algorithm reported as prediction of antiviral drug susceptibility Proprietary test: Sentosa® SQ HIV-1 Genotyping Assay Lab/Manufacturer: Vela Diagnostics USA, Inc
G0432	Infectious agent antibody detection by enzyme immunoassay (EIA) technique, HIV-1 and/or HIV-2, screening
G0433	Infectious agent antibody detection by enzyme-linked immunosorbent assay (ELISA) technique, HIV-1 and/or HIV-2, screening
G0435	Infectious agent antibody detection by rapid antibody test, HIV-1 and/or HIV-2, screening
G0475	HIV antigen/antibody, combination assay, screening
S3645	HIV-1 antibody testing of oral mucosal transudate

Current Procedural Terminology[®] American Medical Association. All Rights reserved.

Reimbursement Policy

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

V. Evidence-based Scientific References

1. CDC. Revised surveillance case definition for HIV infection--United States, 2014. *MMWR Recommendations and reports : Morbidity and mortality weekly report Recommendations and reports*. 2014;63(Rr-03):1-10. April 11, 2014. <https://www.cdc.gov/mmwr/preview/mmwrhtml/rr6303a1.htm>
2. Wood BR. The natural history and clinical features of HIV infection in adults and adolescents. Updated February 15, 2023. <https://www.uptodate.com/contents/the-natural-history-and-clinical-features-of-hiv-infection-in-adults-and-adolescents>
3. Caliendo A. Techniques and interpretation of HIV-1 RNA quantitation. Updated December 9, 2024. <https://www.uptodate.com/contents/techniques-and-interpretation-of-hiv-1-rna-quantitation>
4. FDA. Aptima® HIV-1 Quant Dx Assay. <https://www.fda.gov/media/102425/download>
5. BusinessWire. Aptima HIV-1 Quant Dx Assay Receives Additional FDA Approval for Use as an Aid in the Diagnosis of HIV Infection. 2020. <https://www.businesswire.com/news/home/20201120005242/en/>
6. Sollis KA, Smit PW, Fiscus S, et al. Systematic review of the performance of HIV viral load technologies on plasma samples. *PLoS one*. 2014;9(2):e85869. doi:10.1371/journal.pone.0085869
7. Mor O, Gozlan Y, Wax M, et al. Evaluation of the RealTime HIV-1, Xpert HIV-1, and Aptima HIV-1 Quant Dx Assays in Comparison to the NucliSens EasyQ HIV-1 v2.0 Assay for Quantification of HIV-1 Viral Load. *Journal of clinical microbiology*. Nov 2015;53(11):3458-65. doi:10.1128/jcm.01806-15
8. Swenson LC, Cobb B, Geretti AM, et al. Comparative performances of HIV-1 RNA load assays at low viral load levels: results of an international collaboration. *Journal of clinical microbiology*. Feb 2014;52(2):517-23. doi:10.1128/jcm.02461-13
9. Saag MS, Gandhi RT, Hoy JF, et al. Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2020 Recommendations of the International Antiviral Society—USA Panel. *JAMA*. 2020;324(16):1651-1669. doi:10.1001/jama.2020.17025
10. Saag MS, Holodniy M, Kuritzkes DR, et al. HIV viral load markers in clinical practice. *Nature medicine*. Jun 1996;2(6):625-9. doi:10.1038/nm0696-625

Reimbursement Policy

11. Hughes MD, Johnson VA, Hirsch MS, et al. Monitoring plasma HIV-1 RNA levels in addition to CD4+ lymphocyte count improves assessment of antiretroviral therapeutic response. ACTG 241 Protocol Virology Substudy Team. *Annals of internal medicine*. Jun 15 1997;126(12):929-38. doi:10.7326/0003-4819-126-12-199706150-00001
12. Gottlieb G. Epidemiology, transmission, natural history, and pathogenesis of HIV-2 infection. Updated February 20, 2023. <https://www.uptodate.com/contents/epidemiology-transmission-natural-history-and-pathogenesis-of-hiv-2-infection>
13. Gottlieb G. Clinical manifestations and diagnosis of HIV-2 infection. Updated February 2, 2023. <https://www.uptodate.com/contents/clinical-manifestations-and-diagnosis-of-hiv-2-infection>
14. DHHS. Guidelines for the Use of Antiretroviral Agents in Adults and Adolescents with HIV. Updated September 12, 2024. <https://clinicalinfo.hiv.gov/sites/default/files/guidelines/documents/adult-adolescent-arv/guidelines-adult-adolescent-arv.pdf>
15. Quinn T. Global epidemiology of HIV infection. Updated June 14, 2024. <https://www.uptodate.com/contents/global-epidemiology-of-hiv-infection>
16. Coffin J, Swanstrom R. HIV Pathogenesis: Dynamics and Genetics of Viral Populations and Infected Cells. *Cold Spring Harb Perspect Med*. 2013;3(1)doi:10.1101/cshperspect.a012526
17. Mansky LM, Temin HM. Lower in vivo mutation rate of human immunodeficiency virus type 1 than that predicted from the fidelity of purified reverse transcriptase. *J Virol*. 1995;69(8):5087-94. 189326, Accessed Aug. <https://www.ncbi.nlm.nih.gov/pmc/articles/pmid/7541846/>
18. Kozal M. Interpretation of HIV-1 drug resistance testing. Updated September 17, 2024. <https://www.uptodate.com/contents/interpretation-of-hiv-1-drug-resistance-testing>
19. Kozal M. Overview of HIV-1 drug resistance testing assays. Updated September 17, 2024. <https://www.uptodate.com/contents/overview-of-hiv-1-drug-resistance-testing-assays>
20. ThermoFisher. ViroSeq™ HIV-1 Genotyping System. https://assets.thermofisher.com/TFS-Assets/LSG/brochures/cms_040731.pdf
21. ATCC. ATCC Teams with CDC and Thermo Fisher Scientific on Public Health RT-PCR Assay. https://www.labbulletin.com/articles/ATCC-Teams-CDC-Thermo-Fisher-Scientific-Public-Health-RT-PCR-Assay/categories/20130120_11

Reimbursement Policy

22. LabCorp. Human Immunodeficiency Virus 1 (HIV-1) PhenoSense GT® Plus Integrase (Monogram® Phenotype + Genotype). <https://www.labcorp.com/tests/551920/human-immunodeficiency-virus-1-hiv-1-phenosense-gt-plus-integrase-monogram-phenotype-genotype>
23. Fox ZV, Geretti AM, Kjaer J, et al. The ability of four genotypic interpretation systems to predict virological response to ritonavir-boosted protease inhibitors. *AIDS (London, England)*. Oct 01 2007;21(15):2033-42. doi:10.1097/QAD.0b013e32825a69e4
24. Rosemary A, Chika O, Jonathan O, et al. Genotyping performance evaluation of commercially available HIV-1 drug resistance test. *PloS one*. 2018;13(6):e0198246. doi:10.1371/journal.pone.0198246
25. Braun P, Glass A, Maree L, et al. Multicenter clinical comparative evaluation of Alinity m HIV-1 assay performance. *J Clin Virol*. Aug 2020;129:104530. doi:10.1016/j.jcv.2020.104530
26. Zhang J, Rhee SY, Taylor J, Shafer RW. Comparison of the precision and sensitivity of the Antivirogram and PhenoSense HIV drug susceptibility assays. *Journal of acquired immune deficiency syndromes (1999)*. Apr 1 2005;38(4):439-44. doi:10.1097/01.qai.0000147526.64863.53
27. Hopkins M, Hau S, Tiernan C, et al. Comparative performance of the new Aptima HIV-1 Quant Dx assay with three commercial PCR-based HIV-1 RNA quantitation assays. *Journal of Clinical Virology*. 2015/08/01/ 2015;69:56-62. doi:10.1016/j.jcv.2015.05.020
28. Sempa JB, Dushoff J, Daniels MJ, et al. Reevaluating Cumulative HIV-1 Viral Load as a Prognostic Predictor: Predicting Opportunistic Infection Incidence and Mortality in a Ugandan Cohort. *American journal of epidemiology*. Jul 1 2016;184(1):67-77. doi:10.1093/aje/kwv303
29. Shen C, Yu X, Harrison RW, Weber IT. Automated prediction of HIV drug resistance from genotype data. *BMC bioinformatics*. Aug 31 2016;17 Suppl 8:278. doi:10.1186/s12859-016-1114-6
30. Lindman J, Hønge BL, Kjerulff B, et al. Performance of Bio-Rad HIV-1/2 Confirmatory Assay in HIV-1, HIV-2 and HIV-1/2 dually reactive patients - comparison with INNO-LIA and immunocomb discriminatory assays. *J Virol Methods*. Jun 2019;268:42-47. doi:10.1016/j.jviromet.2019.03.005
31. Avram CM, Greiner KS, Tilden E, Caughey AB. Point-of-care HIV viral load in pregnant women without prenatal care: a cost-effectiveness analysis. *Am J Obstet Gynecol*. Sep 2019;221(3):265.e1-265.e9. doi:10.1016/j.ajog.2019.06.021

Reimbursement Policy

32. Raymond S, Nicot F, Abravanel F, et al. Performance evaluation of the Vela Dx Sentosa next-generation sequencing system for HIV-1 DNA genotypic resistance. *Journal of Clinical Virology*. 2020/01/01/ 2020;122:104229. doi:10.1016/j.jcv.2019.104229
33. Fogel JM, Bonsall D, Cummings V, et al. Performance of a high-throughput next-generation sequencing method for analysis of HIV drug resistance and viral load. *Journal of Antimicrobial Chemotherapy*. 2020;75(12):3510-3516. doi:10.1093/jac/dkaa352
34. Pröll J, Paar C, Taylor N, et al. New aspects of the Virus Life Cycle and Clinical Utility of Next Generation Sequencing based HIV-1 Resistance Testing in the Genomic, the Proviral and the Viral Reservoir of Peripheral Blood Mononuclear Cells. *Curr HIV Res*. Mar 24 2022;doi:10.2174/1570162x20666220324111418
35. Ehret R, Harb K, Breuer S, Obermeier M. Performance assessment of the new Xpert® HIV-1 viral load XC assay for quantification of HIV-1 viral loads. *J Clin Virol*. Apr 2022;149:105127. doi:10.1016/j.jcv.2022.105127
36. DHHS. Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection. Updated December 19, 2024. <https://clinicalinfo.hiv.gov/en/guidelines/pediatric-arv/whats-new-guidelines>
37. DHHS. Recommendations for the Use of Antiretroviral Drugs in Pregnant Women with HIV Infection and Interventions to Reduce Perinatal HIV Transmission in the United States. Updated December 19, 2024. <https://clinicalinfo.hiv.gov/en/guidelines/perinatal/whats-new-guidelines>
38. Gandhi RT, Bedimo R, Hoy JF, et al. Antiretroviral Drugs for Treatment and Prevention of HIV Infection in Adults: 2022 Recommendations of the International Antiviral Society–USA Panel. *JAMA*. 2022;doi:10.1001/jama.2022.22246
39. Thompson MA, Horberg MA, Agwu AL, et al. Primary Care Guidance for Persons With Human Immunodeficiency Virus: 2020 Update by the HIV Medicine Association of the Infectious Diseases Society of America. *Clinical Infectious Diseases*. 2020;doi:10.1093/cid/ciaa1391
40. ACOG. ACOG Committee Opinion no 596: Committee on Gynecologic Practice: Routine human immunodeficiency virus screening. *Obstet Gynecol*. May 2014;123(5):1137-1139. doi:10.1097/01.AOG.0000446828.64137.50
41. ACOG. ACOG Committee Opinion No. 751: Labor and Delivery Management of Women With Human Immunodeficiency Virus Infection. *Obstet Gynecol*. Sep 2018;132(3):e131-e137. doi:10.1097/aog.0000000000002820

Reimbursement Policy

42. Gibson KS, Toner LE. Society for Maternal-Fetal Medicine Special Statement: Updated checklists for pregnancy management in persons with HIV. *American Journal of Obstetrics & Gynecology*. 2020;223(5):B6-B11. doi:10.1016/j.ajog.2020.08.064
43. BHIVA. BHIVA guidelines for the routine investigation and monitoring of adult HIV-1-positive individuals (2019 interim update). <https://www.bhiva.org/file/DqZbRxfzLYtLg/Monitoring-Guidelines.pdf>
44. BHIVA. British HIV Association guidelines for the management of HIV-2 2021. 2021. <https://www.bhiva.org/file/615ee3de98539/BHIVA-guidelines-for-the-management-of-HIV-2.pdf>
45. EACS. European AIDS Clinical Society Guidelines Version 11.1 October 2022. https://www.eacsociety.org/media/guidelines-11.1_final_09-10.pdf
46. Hsu KK, Rakhmanina NY. Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis. *Pediatrics*. Jan 1 2022;149(1)doi:10.1542/peds.2021-055207
47. AAP. Bright Futures/AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule). <https://www.aap.org/periodicityschedule>
48. CDC. Getting Tested for HIV. <https://www.cdc.gov/hiv/testing/index.html>
49. CDC. HIV Infection. <https://www.cdc.gov/immigrant-refugee-health/hcp/domestic-guidance/hiv-infection.html>
50. USPSTF. Screening for HIV Infection: US Preventive Services Task Force Recommendation Statement. *JAMA*. 2019;321(23):2326-2336. doi:10.1001/jama.2019.6587
51. CDC. HIV Preventative Services Coverage. <https://www.cdc.gov/high-quality-care/hcp/resources/hiv-preventative-service-coverage.html>
52. FDA. Abbott RealTime HIV-1 <https://www.fda.gov/media/73278/download>
53. FDA. COBAS® AmpliPrep/COBAS® TaqMan® HIV-1 Test, Summary of Safety and Effectiveness <https://www.fda.gov/media/73824/download>
54. FDA. Geenius™ HIV 1/2 Supplemental Assay. <https://www.fda.gov/media/130312/download>
55. FDA. BioPlex 2200 HIV Ag-Ab Assay. <https://www.fda.gov/media/92862/download>
56. FDA. Alinity m HIV-1. <https://www.fda.gov/vaccines-blood-biologics/alinity-m-hiv-1>