

Reimbursement Policy

Urinary Tumor Markers for Bladder Cancer

[POLICY DESCRIPTION](#) | [INDICATIONS AND/OR LIMITATIONS OF COVERAGE](#) | [APPLICABLE STATE AND FEDERAL REGULATIONS](#) | [APPLICABLE CPT/HCPCS PROCEDURE CODES](#) | [EVIDENCE-BASED SCIENTIFIC REFERENCES](#) | [REVISION HISTORY](#)

I. Policy Description

Bladder cancer is defined as a malignancy that develops from the tissues of the bladder. It is the most common cancer of the urinary system. The cancer typically arises from the urothelium, although it may originate in other locations such as the ureter or urethra.¹

Tumor biomarkers are proteins detected in the blood, urine, or other body fluids that are produced by the tumor itself or in response to it. Urinary tumor markers may be used to help detect, diagnose, and manage some types of cancer including bladder cancer.²

II. Indications and/or Limitations of Coverage

Application of coverage criteria is dependent upon an individual's benefit coverage at the time of the request. Specifications pertaining to Medicare and Medicaid can be found in the "Applicable State and Federal Regulations" section of this policy document.

- 1) Urinary biomarkers (bladder tumor antigen (BTA) test, nuclear matrix protein (NMP22) test, or fluorescence in situ hybridization (FISH) UroVysion Bladder Cancer test) **MEET COVERAGE CRITERIA** in any of the following situations:
 - a) As an adjunct in the diagnostic exclusion of bladder cancer for individuals who have an atypical or equivocal cytology.
 - b) As an adjunct in the monitoring of high-risk, non-muscle invasive bladder cancer.
- 2) As an adjunct to cystoscopy or cytology in the monitoring of individuals with bladder cancer, the use of fluorescence immunocytology (ImmunoCyt/uCyt) **MEETS COVERAGE CRITERIA**.

The following does not meet coverage criteria due to a lack of available published scientific literature confirming that the test(s) is/are required and beneficial for the diagnosis and treatment of an individual's illness.

- 3) For the evaluation of hematuria, to screen for bladder cancer in asymptomatic individuals, to diagnose bladder cancer in symptomatic individuals, or for any other indication not discussed above, the following tests **DO NOT MEET COVERAGE CRITERIA**:
 - a) Urinary biomarkers (bladder tumor antigen (BTA) test, nuclear matrix protein (NMP22) test, or fluorescence in situ hybridization (FISH) UroVysion Bladder Cancer test).

Reimbursement Policy

- b) Fluorescence immunocytology (ImmunoCyt/uCyt).
- 4) Any other urinary tumor markers for bladder cancer not mentioned above **DO NOT MEET COVERAGE CRITERIA.**

III. Applicable State and Federal Regulations

DISCLAIMER: If there is a conflict between this policy and any relevant, applicable government policy for a particular member [e.g., Local Coverage Determinations (LCDs) or National Coverage Determinations (NCDs) for Medicare and/or state coverage for Medicaid], then the government policy will be used to make the determination. For the most up-to-date Medicare policies and coverage, please visit the Medicare search website: <http://www.cms.gov/medicare-coverage-database/search.aspx>. For the most up-to-date Medicaid policies and coverage, visit the applicable state Medicaid website.

Food and Drug Administration

Many labs have developed specific tests that they must validate and perform in house. These laboratory-developed tests (LDTs) are regulated by the Centers for Medicare and Medicaid (CMS) as high-complexity tests under the Clinical Laboratory Improvement Amendments of 1988 (CLIA '88). LDTs are not approved or cleared by the U. S. Food and Drug Administration; however, FDA clearance or approval is not currently required for clinical use.

On April 16, 1997, the FDA approved the *Bard BTA stat™ Test*, created by Bard Diagnostic Sciences Inc. From the FDA site: “the BTA stat test is an *in vitro* diagnostic immunoassay indicated for the qualitative detection of bladder tumor associated antigen in urine of persons diagnosed with bladder cancer. This test is indicated for use as an aid in the management of bladder cancer patients in conjunction with cystoscopy.”⁶³

On April 15, 1998, the FDA approved the *BTA TRAK™ Test*, created by Bard Diagnostic Sciences Inc. From the FDA site: “the BTA TRAK test is an *in vitro* diagnostic immunoassay indicated for the quantitative detection of bladder tumor associated antigen in human urine. This test is indicated for use as an aid in the management of bladder cancer patients in conjunction with cystoscopy.”⁶⁴

On July 2, 1996, the FDA approved the *MATRITECH NMP22™ TEST KIT*, created by Alere Scarborough Inc. From the FDA site: “The Matritech NMP22 Test Kit is an enzyme immunoassay (EIA) for the *in vitro* quantitative determination of nuclear matrix protein NMP22 in stabilized voided urine.”⁶⁵

On July 30, 2002, the FDA approved the *NMP22 BladderChek*, created by Matritech Inc. From the FDA site: “The Matritech NMP22 BladderChek Test is indicated for professional and prescription home use as an aid in monitoring bladder cancer patients, in conjunction with standard diagnostic procedures.” This assay is qualitative.⁶⁶

On January 24, 2005, the FDA approved the *UROVYSION BLADDER CANCER KIT*. From the FDA site: “The UroVysion Bladder Cancer Kit (UroVysion Kit) is designed to detect aneuploidy for chromosomes 3, 7, 17, and loss of the 9p21 locus via fluorescence *in situ* hybridization (FISH) in urine specimens from persons with hematuria suspected of having bladder cancer.”⁶⁷

Reimbursement Policy

On February 23, 2000, the FDA approved the *ImmunoCyt*, created by Diagnostics Inc. From the FDA site: “ImmunoCyt is a qualitative direct immunofluorescence assay intended for use in conjunction with cytology to increase overall sensitivity for the detection of tumor cells exfoliated in the urine of patients previously diagnosed with bladder cancer. ImmunoCyt is indicated for use as an aid in the management of bladder cancer in conjunction with urinary cytology and cystoscopy.”⁶⁸

All of the FDA-approved tests apart from ImmunoCyt are approved for both diagnosis and surveillance of bladder cancer whereas ImmunoCyt is only approved for surveillance.⁶⁹

IV. Applicable CPT/HCPCS Procedure Codes

CPT	Code Description
86294	Immunoassay for tumor antigen, qualitative or semiquantitative (eg, bladder tumor antigen)
86316	Immunoassay for tumor antigen, other antigen, quantitative (eg, CA 50, 72-4, 549), each
86386	Nuclear Matrix Protein 22 (NMP22), qualitative
88120	Cytopathology, in situ hybridization (eg, FISH), urinary tract specimen with morphometric analysis, 3-5 molecular probes, each specimen; manual
88121	Cytopathology, in situ hybridization (eg, FISH), urinary tract specimen with morphometric analysis, 3-5 molecular probes, each specimen; using computer-assisted technology
88346	Immunofluorescence, per specimen; initial single antibody stain procedure
88350	Immunofluorescence, per specimen; each additional single antibody stain procedure (List separately in addition to code for primary procedure)
0365U	Oncology (bladder), analysis of 10 protein biomarkers (A1AT, ANG, APOE, CA9, IL8, MMP9, MMP10, PAI1, SDC1 and VEGFA), by immunoassays, urine, diagnostic algorithm, including patient's age, race and gender, reported as a probability of harboring urothelial bladder cancer Proprietary test: Oncuria® Detect Lab/Manufacturer: DiaCarta Clinical Lab
0366U	Oncology (bladder), analysis of 10 protein biomarkers (A1AT, ANG, APOE, CA9, IL8, MMP9, MMP10, PAI1, SDC1 and VEGFA) by immunoassays, urine, algorithm reported as a probability of recurrent bladder cancer Proprietary test: Oncuria® Monitor Lab/Manufacturer: DiaCarta Clinical Lab
0367U	Oncology (bladder), analysis of 10 protein biomarkers (A1AT, ANG, APOE, CA9, IL8, MMP9, MMP10, PAI1, SDC1 and VEGFA) by immunoassays, urine, diagnostic algorithm reported as a risk score for probability of rapid recurrence of recurrent or persistent cancer following transurethral resection Proprietary test: Oncuria® Predict Lab/Manufacturer: DiaCarta Clinical Lab

Current Procedural Terminology© American Medical Association. All Rights reserved.

Procedure codes appearing in Medical Policy documents are included only as a general reference tool for each policy. They may not be all-inclusive.

Reimbursement Policy

V. Evidence-based Scientific References

1. Lerner SP, Raghavan, Derek. Overview of the initial approach and management of urothelial bladder cancer. Updated July 29, 2024. <https://www.uptodate.com/contents/overview-of-the-initial-approach-and-management-of-urothelial-bladder-cancer>
2. Hottinger AF, Hormigo A. Serum Biomarkers. In: Schwab M, ed. *Encyclopedia of Cancer*. Springer Berlin Heidelberg; 2011:3390-3394.
3. ACS. Key Statistics for Bladder Cancer. Updated March 12, 2024. <https://www.cancer.org/cancer/types/bladder-cancer/about/key-statistics.html>
4. NCCN. NCCN Guidelines Version 5.2024 Bladder Cancer. Updated October 28, 2024. https://www.nccn.org/professionals/physician_gls/pdf/bladder.pdf
5. DeGeorge KC, Holt HR, Hodges SC. Bladder Cancer: Diagnosis and Treatment. 2017;(1532-0650 (Electronic))
6. Perazalla M. Evaluation of hematuria in adults. Updated December 18, 2024. <https://www.uptodate.com/contents/evaluation-of-hematuria-in-adults>
7. Lotan Y, Choueiri T. Clinical presentation, diagnosis, and staging of bladder cancer. Updated February 25, 2025. <https://www.uptodate.com/contents/clinical-presentation-diagnosis-and-staging-of-bladder-cancer>
8. Sutton AJ, Lamont JV, Evans RM, et al. An early analysis of the cost-effectiveness of a diagnostic classifier for risk stratification of haematuria patients (DCRSHP) compared to flexible cystoscopy in the diagnosis of bladder cancer. *PloS one*. 2018;13(8):e0202796. doi:10.1371/journal.pone.0202796
9. Kaufman DS, Shipley WU, Feldman AS. Bladder cancer. *Lancet (London, England)*. Jul 18 2009;374(9685):239-49. doi:10.1016/s0140-6736(09)60491-8
10. Chou R, Dana T. Screening adults for bladder cancer: A review of the evidence for the u.s. preventive services task force. *Annals of Internal Medicine*. 2010;153(7):461-468. doi:10.7326/0003-4819-153-7-201010050-00009
11. Mitra A, Birkman M, Penson D, Cote R. Urine biomarkers for the detection of urothelial (transitional cell) carcinoma of the bladder Updated January 18, 2024. <https://www.uptodate.com/contents/urine-biomarkers-for-the-detection-of-urothelial-transitional-cell-carcinoma-of-the-bladder>
12. Lopez-Beltran A, Cheng L, Gevaert T, et al. Current and emerging bladder cancer biomarkers with an emphasis on urine biomarkers. *Expert Rev Mol Diagn*. Dec 11 2019;1-13. doi:10.1080/14737159.2020.1699791
13. D'Costa JJ, Goldsmith JC, Wilson JS, Bryan RT, Ward DG. A Systematic Review of the Diagnostic and Prognostic Value of Urinary Protein Biomarkers in Urothelial Bladder Cancer. *Bladder cancer (Amsterdam, Netherlands)*. 2016;2(3):301-317. doi:10.3233/blc-160054
14. Lotan Y, Elias K, Svatek RS, et al. Bladder cancer screening in a high risk asymptomatic population using a point of care urine based protein tumor marker. *The Journal of urology*. Jul 2009;182(1):52-7; discussion 58. doi:10.1016/j.juro.2009.02.142
15. Li HT, Duymich CE, Weisenberger DJ, Liang G. Genetic and Epigenetic Alterations in Bladder Cancer. *Int Neurol J*. Nov 2016;20(Suppl 2):S84-94. doi:10.5213/inj.1632752.376
16. Mahnert B, Tauber S, Kriegmair M, et al. BTA-TRAK--a useful diagnostic tool in urinary bladder cancer? *Anticancer research*. Jul-Aug 1999;19(4a):2615-9.
17. Grossman H, Messing E, Soloway M, et al. Detection of bladder cancer using a point-of-care proteomic assay. *JAMA*. 2005;293(7):810-816. doi:10.1001/jama.293.7.810

Reimbursement Policy

18. Zuiverloon TCM, de Jong FC, Theodorescu D. Clinical Decision Making in Surveillance of Non-Muscle-Invasive Bladder Cancer: The Evolving Roles of Urinary Cytology and Molecular Markers. *Oncology (Williston Park, NY)*. Dec 15 2017;31(12):855-62.
19. Abbott. ALERE NMP22® BLADDERCHEK®. <https://www.globalpointofcare.abbott/en/product-details/nmp22-bladderchek.html>
20. Pangea. Features of Bladder CARE™. <https://pangealab.com/bladder-care/>
21. Pangea. Pangea® Laboratory to License Bladder CARE™ Technology from Zymo Research. <https://www.prnewswire.com/news-releases/pangea-laboratory-to-license-bladder-care-technology-from-zymo-research-300800622.html>
22. Nucleix. BLADDER EPICHECK <https://www.nucleix.com/bladder-epicheck/>
23. D'Andrea D, Soria F, Zehetmayer S, et al. Diagnostic accuracy, clinical utility and influence on decision-making of a methylation urine biomarker test in the surveillance of non-muscle-invasive bladder cancer. *BJU Int*. Jun 2019;123(6):959-967. doi:10.1111/bju.14673
24. AroCell. Bladder Cancer Detection. <https://arocell.com/products/oncology/ubc-rapid/>
25. Ecke TH, Weiß S, Stephan C, et al. UBC(®) Rapid Test-A Urinary Point-of-Care (POC) Assay for Diagnosis of Bladder Cancer with a focus on Non-Muscle Invasive High-Grade Tumors: Results of a Multicenter-Study. *Int J Mol Sci*. Dec 2 2018;19(12)doi:10.3390/ijms19123841
26. NICE. URO17 for detecting bladder cancer. Updated February 4, 2021. <https://www.nice.org.uk/advice/mib250/chapter/The-technology>
27. KdX Diagnostics. URO17® A Better test For Better Health. <https://kdxdiagnostics.com/uro17/>
28. Vasdev, Hampson A, Agarwal S, et al. The role of URO17™ biomarker to enhance diagnosis of urothelial cancer in new hematuria patients-First European Data. *BJUI Compass*. Jan 2021;2(1):46-52. doi:10.1002/bco2.50
29. Nonagen Bioscience. Bladder Cancer. <https://www.nonagen.com/products>
30. Pichler R, Fritz J, Tulchiner G, et al. Increased accuracy of a novel mRNA-based urine test for bladder cancer surveillance. *BJU Int*. Jan 2018;121(1):29-37. doi:10.1111/bju.14019
31. D'Elia C, Folchini DM, Mian C, et al. Diagnostic value of Xpert(®) Bladder Cancer Monitor in the follow-up of patients affected by non-muscle invasive bladder cancer: an update. *Ther Adv Urol*. Jan-Dec 2021;13:1756287221997183. doi:10.1177/1756287221997183
32. Piao XM, Jeong P, Kim YH, et al. Urinary cell-free microRNA biomarker could discriminate bladder cancer from benign hematuria. *Int J Cancer*. Jan 15 2019;144(2):380-388. doi:10.1002/ijc.31849
33. Fantony JJ, Longo TA, Gopalakrishna A, et al. Urinary NID2 and TWIST1 methylation to augment conventional urine cytology for the detection of bladder cancer. *Cancer Biomark*. 2017;18(4):381-387. doi:10.3233/cbm-160261
34. Soubra A, Risk MC. Diagnostics techniques in nonmuscle invasive bladder cancer. *Indian journal of urology : IJU : journal of the Urological Society of India*. Oct-Dec 2015;31(4):283-8. doi:10.4103/0970-1591.166449
35. Lotan Y, Roehrborn CG. Sensitivity and specificity of commonly available bladder tumor markers versus cytology: results of a comprehensive literature review and meta-analyses. *Urology*. 2003;61(1):109-118. doi:10.1016/S0090-4295(02)02136-2
36. Breen V, Kasabov N, Kamat AM, et al. A holistic comparative analysis of diagnostic tests for urothelial carcinoma: a study of Cxbladder Detect, UroVysion® FISH, NMP22® and cytology based on imputation of multiple datasets. *BMC medical research methodology*. 2015;15:45-45. doi:10.1186/s12874-015-0036-8

Reimbursement Policy

37. Sathianathen NJ, Butaney M, Weight CJ, Kumar R, Konety BR. Urinary Biomarkers in the Evaluation of Primary Hematuria: A Systematic Review and Meta-Analysis. *Bladder cancer (Amsterdam, Netherlands)*. 2018;4(4):353-63.
38. Chou R, Gore JL, Buckley D, et al. Urinary Biomarkers for Diagnosis of Bladder Cancer: A Systematic Review and Meta-analysis. *Ann Intern Med*. Dec 15 2015;163(12):922-31. doi:10.7326/m15-0997
39. Dudley JC, Schroers-Martin J, Lazzareschi DV, et al. Detection and Surveillance of Bladder Cancer Using Urine Tumor DNA. *Cancer Discov*. Apr 2019;9(4):500-509. doi:10.1158/2159-8290.Cd-18-0825
40. Hirasawa Y, Pagano I, Chen R, et al. Diagnostic performance of Oncuria™, a urinalysis test for bladder cancer. *Journal of Translational Medicine*. 2021/04/06 2021;19(1):141. doi:10.1186/s12967-021-02796-4
41. Schmitz-Dräger BJ, Droller M, Lokeshwar VB, et al. Molecular Markers for Bladder Cancer Screening, Early Diagnosis, and Surveillance: The WHO/ICUD Consensus. *Urologia Internationalis*. 2015;94(1):1-24. doi:10.1159/000369357
42. Halpern JA, Chughtai B, Ghomrawi H. Cost-effectiveness of Common Diagnostic Approaches for Evaluation of Asymptomatic Microscopic Hematuria. *JAMA internal medicine*. Jun 2017;177(6):800-807. doi:10.1001/jamainternmed.2017.0739
43. Landaas EJ, Eckel AM, Wright JL, Baird GS, Hansen RN, Sullivan SD. Application of Health Technology Assessment (HTA) to Evaluate New Laboratory Tests in a Health System: A Case Study of Bladder Cancer Testing. *Acad Pathol*. Jan-Dec 2020;7:2374289520968225. doi:10.1177/2374289520968225
44. Meleth S, Reeder-Hayes K, Ashok M, et al. AHRQ Technology Assessments. *Technology Assessment of Molecular Pathology Testing for the Estimation of Prognosis for Common Cancers*. Agency for Healthcare Research and Quality (US); 2014.
45. Tan WS, Tan WP, Tan MY, et al. Novel urinary biomarkers for the detection of bladder cancer: A systematic review. *Cancer Treat Rev*. Sep 2018;69:39-52. doi:10.1016/j.ctrv.2018.05.012
46. Mossanen M, Wang Y, Szymaniak J, et al. Evaluating the cost of surveillance for non-muscle-invasive bladder cancer: an analysis based on risk categories. *World J Urol*. Oct 2019;37(10):2059-2065. doi:10.1007/s00345-018-2550-x
47. Vasdev N, Hampson A, Agarwal S, et al. The role of URO17™ biomarker to enhance diagnosis of urothelial cancer in new hematuria patients—First European Data. *BJUI Compass*. 2021;2(1):46-52. doi:10.1002/bco2.50
48. Davis R, Jones JS, Barocas DA, et al. Diagnosis, evaluation and follow-up of asymptomatic microhematuria (AMH) in adults: AUA guideline. *The Journal of urology*. Dec 2012;188(6 Suppl):2473-81. doi:10.1016/j.juro.2012.09.078
49. Barocas DA, Boorjian SA, Alvarez RD, et al. Microhematuria: AUA/SUFU Guideline. *The Journal of urology*. Oct 2020;204(4):778-786. doi:10.1097/ju.0000000000001297
50. Chang SS, Boorjian SA, Chou R, et al. Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Guideline. *The Journal of urology*. Oct 2020;196(4):1021-9. doi:10.1016/j.juro.2016.06.049
51. Holzbeierlein J BB, Buckley DI, et al. Diagnosis and Treatment of Non-Muscle Invasive Bladder Cancer: AUA/SUO Guideline: 2024 Amendment. *Journal of Urology [Internet]*. 2024 Apr 1 2024;

Reimbursement Policy

52. Holzbeierlein J BB, Buckley DI, et al. Treatment of non-metastatic muscle-invasive bladder cancer: AUA/ASCO/SUO guideline (2017; amended 2020, 2024). *J Urol*. 2024;
53. Chang SS, Bochner BH, Chou R, et al. Treatment of Non-Metastatic Muscle-Invasive Bladder Cancer: AUA/ASCO/ASTRO/SUO Guideline. *The Journal of urology*. Sep 2017;198(3):552-559. doi:10.1016/j.juro.2017.04.086
54. Moyer VA. Screening for bladder cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. Aug 16 2011;155(4):246-51. doi:10.7326/0003-4819-155-4-201108160-00008
55. USPSTF. Bladder Cancer in Adults: Screening. <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/bladder-cancer-in-adults-screening>
56. Monteiro LL, Witjes JA, Agarwal PK, et al. ICUD-SIU International Consultation on Bladder Cancer 2017: management of non-muscle invasive bladder cancer. 2018:1-10.
57. NCI. Bladder and Other Urothelial Cancers Screening (PDQ®)—Health Professional Version. U.S. Department of Health and Human Services. Accessed 12/12/2024, <https://www.cancer.gov/types/bladder/hp/bladder-screening-pdq>
58. Witjes JA, Bruins HM, Cathomas R, et al. Muscle-invasive and Metastatic Bladder Cancer. <https://doi.org/10.1016/j.eururo.2023.08.016>
59. Babjuk M, Burger M, Capoun O, et al. European Association of Urology Guidelines on Non-muscle-invasive Bladder Cancer (Ta, T1, and Carcinoma in Situ). *European Urology*. 2022;81(1):75-94. doi:10.1016/j.eururo.2021.08.010
60. EAU. EAU Guidelines on Non-muscle Invasive Bladder Cancer. <https://uroweb.org/guidelines/non-muscle-invasive-bladder-cancer>
61. Bhindi B, Kool R, Kulkarni GS, et al. Canadian Urological Association guideline on the management of non-muscle-invasive bladder cancer – Full-text. *Can Urol Assoc J* 2021;
62. Stecca C, Chowdhury D, Blais N, et al. 2024 CUA-GUMOC Expert Report: Management of unresectable locally advanced and metastatic urothelial carcinoma. *Can Urol Assoc J*. 2024;18(12):379-90
63. FDA. 510(k) Summary Information. https://www.accessdata.fda.gov/cdrh_docs/pdf/K964151.pdf
64. FDA. 510(k) Summary Information. https://www.accessdata.fda.gov/cdrh_docs/pdf/K971402.pdf
65. FDA. Premarket Approval. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P940035>
66. FDA. 510(k) Summary. https://www.accessdata.fda.gov/cdrh_docs/pdf2/K021231.pdf
67. FDA. Premarket Approval. <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P030052>
68. FDA. 510(k) number: K994356. https://www.accessdata.fda.gov/cdrh_docs/pdf/K994356.pdf
69. Darwiche F, Parekh DJ, Gonzalgo ML. Biomarkers for non-muscle invasive bladder cancer: Current tests and future promise. *Indian journal of urology : IJU : journal of the Urological Society of India*. Oct-Dec 2015;31(4):273-82. doi:10.4103/0970-1591.166448

Reimbursement Policy

VI. Revision History

Revision Date	Summary of Changes
12/03/2025	Off-cycle coding modification: No changes to policy outside of coding section table. Removed CPT code 0012M, 0013M, 0363U, 0420U, 0452U, 0465U, 0549U (now managed in GTM policy G2054); Removed CPT code 0613U (mapped to policy during initial 1/1/2026 PLA review but changes to policy required moving 0613U to GTM policy G2054)